

RESEARCH ARTICLE

Open Access

Antibacterial activities of selected Cameroonian spices and their synergistic effects with antibiotics against multidrug-resistant phenotypes

Aimé G Fankam¹, Victor Kuete^{1*}, Igor K Voukeng¹, Jules R Kuate¹ and Jean-Marie Pages²

Abstract

Background: The emergence of multi-drug resistant (MDR) phenotypes is a major public health problem today in the treatment of bacterial infections. The present study was designed to evaluate the antibacterial activities of the methanol extracts of eleven Cameroonian spices on a panel of twenty nine Gram negative bacteria including MDR strains.

Methods: The phytochemical analysis of the extracts was carried out by standard tests meanwhile the liquid micro-broth dilution was used for all antimicrobial assays.

Results: Phytochemical analysis showed the presence of alkaloids, phenols and tannins in all plants extracts. The results of the antibacterial assays indicated that all tested extracts exert antibacterial activities, with the minimum inhibitory concentration (MIC) values varying from 32 to 1024 µg/ml. The extracts from *Dichrostachys glomerata*, *Beilschmiedia cinnamomea*, *Aframomum citratum*, *Piper capense*, *Echinops giganteus*, *Fagara xanthoxyloides* and *Olax subscorpioidea* were the most active. In the presence of efflux pump inhibitor, PAβN, the activity of the extract from *D. glomerata* significantly increased on 69.2% of the tested MDR bacteria. At MIC/5, synergistic effects were noted with the extract of *D. glomerata* on 75% of the tested bacteria for chloramphenicol (CHL), tetracycline (TET) and norfloxacin (NOR). With *B. cinnamomea* synergy were observed on 62.5% of the studied MDR bacteria with CHL, cefepime (FEP), NOR and ciprofloxacin (CIP) and 75% with erythromycin (ERY).

Conclusion: The overall results provide information for the possible use of the studied extracts of the spices in the control of bacterial infections involving MDR phenotypes.

Background

The emergence of MDR phenotypes is a major public health problem today in the treatment of bacterial infections. The multi-drug resistance of Gram negative bacteria is a major cause of morbidity and mortality in health care services [1]. The activation of bacterial efflux pumps also plays an important role in the appearance of resistance to antibiotics [2]. The real challenge for scientists worldwide today, is to continuously find new drugs to combat resistant microorganisms, or compounds which are able to inhibit the resistance mechanisms of pathogens, therefore restoring the activity of antibiotics. Medicinal plants are rich in compounds which may be potential natural drugs

and serve as alternative, less expensive and safe antimicrobials for the treatment of common ailments. Plant drugs are widely used in Africa for the treatment of many ailments and constitute the first health recourse for about 80% of the population [3]. A number of pharmaceutical products in current use worldwide are derived from plants [4]. In Cameroon, many medicinal plants including spices are used as herbal medicines. The present work was therefore designed to investigate the antibacterial potential against MDR bacteria of some of the commonly used medicinal spices in Cameroon such as *Fagara xanthoxyloides* Watern., *Dichrostachys glomerata* (Forsk) Chuov., *Olax subscorpioidea* Oliv., *Solanum melongena* L. Var inerme D.C Hiern, *Piper capense* Lin.f, *Xylopi aethiopica* Dunal A. Rich., *Aframomum citratum* (Pereira). Schum, *Scorodophloeus zenkeri* Harms., *Beilschmiedia cinnamomea* (Stapf) Robyns & Wilczek, *Echinops giganteus* A. Rich

* Correspondence: kuetevictor@yahoo.fr

¹Department of Biochemistry, Faculty of science, University of Dschang, Cameroon

Full list of author information is available at the end of the article

and *Mondia whitei* (Hook F). Skell. This study was also extended to the evaluation of the potencies of the above plant extracts to increase the activity of some antibiotics on MDR bacteria. The role of bacterial efflux pumps in resistance to the extracts was also studied.

Methods

Plant materials and extraction

The eleven edible spices used in this work were purchased from Dschang local market, West Region of Cameroon in January 2010. The collected spices materials were: the fruits of *Fagara xanthoxyloides*, *Dichrostachys glomerata*, *Olox subscorpioidea*, *Solanum melongena*, *Piper capense* and *Xylopiya aethiopica*, the bark of *Aframomum citratum*, *Scorodophloeus zenkeri*, *Beilschmiedia cinnamomea* and the roots of *Echinops giganteus* and *Mondia whitei*. The plants were identified by Mr. Fulbert Tadjouteu of the National herbarium (Yaoundé, Cameroon) where voucher specimens were deposited under the reference numbers (Table 1).

The air dried and powdered sample (1 kg) from each spice was extracted with methanol (MeOH) for 48 h at room temperature. The extract was then concentrated under reduced pressure to give residues which constituted the crude extracts. They were then kept at 4°C until further use.

Preliminary phytochemical investigations

The major classes of secondary metabolites; alkaloids, anthocyanins, anthraquinones, flavonoids, phenols, saponins, tannins, steroids and triterpenes were screened according to the common phytochemical methods described by Harborne [5] with some modifications. Briefly, for alkaloids (5 mg plant extract in 10 ml methanol); a portion of 2 ml extract + 1% HCl + steam, 1 ml filtrate + 6 drops of Mayor's reagents/Wagner's reagent/ Dragendroff reagent; creamish precipitate/brownish-red precipitate/orange precipitate indicated the presence of respective alkaloids. For tannins (5 mg plant extract in 10 ml distilled water); a portion of 2 ml + 2 ml FeCl₃; blue-black precipitate indicated the presence of tannins. For saponins (frothing test: 0.5 ml filtrate + 5 ml distilled water); frothing persistence indicated presence of saponins. For steroids and triterpenoids (Liebermann-Burchard reaction: 5 mg plant extract in 10 ml chloroform, filtered); a 2 ml filtrate + 2 ml acetic anhydride + conc. H₂SO₄. Blue-green ring or pink-purple indicated the presence of steroids or triterpenoids. For flavonoids (5 mg plant extract in 10 ml methanol); a portion of 2 ml + conc. HCl + magnesium; ribbon pink-tomato red color indicated the presence of flavonoids. For anthocyanins (5 mg plant extract in 10 ml methanol); a portion 2 ml + 1% HCl + heating; orange color indicated the presence of anthocyanins. For anthraquinones (5 mg plant extract in

10 ml methanol); a portion of 2 ml + 2 ml ether-chloroform 1:1 v/v + 4 ml NaOH 10% (w/v); red color indicated the presence of anthraquinones. For phenols (5 mg plant material in 10 ml methanol); a portion of 2 ml + 2 ml FeCl₃; violet-blue or greenish color indicated the presence of phenols.

Chemicals for antimicrobial assays

Tetracycline (TET), cefepime (FEP), streptomycin (STR), ciprofloxacin (CIP), norfloxacin (NOR), chloramphenicol (CHL), cloxacillin (CLX), ampicillin (AMP), erythromycin (ERY), kanamycin (KAN) (Sigma-Aldrich, St Quentin Fallavier, France) were used as reference antibiotics. *p*-Iodonitrotetrazolium chloride (INT) and phenylalanine arginine β -naphthylamide (PA β N) were used as microbial growth indicator and efflux pumps inhibitor respectively.

Bacterial strains and culture media

The studied microorganisms included reference (from the American Type Culture Collection) and clinical (Laboratory collection) strains of *Providencia stuartii*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Escherichia coli*, *Enterobacter aerogenes* and *Enterobacter cloacae* (Table 2). They were maintained on agar slant at 4°C and sub-cultured on a fresh appropriate agar plates 24 h prior to any antimicrobial test. Mueller Hinton Agar was used for the activation of bacteria. The Mueller Hinton Broth (MHB) was used for the MIC determinations.

Bacterial susceptibility determinations

The respective MICs of samples on the studied bacteria were determined using rapid INT colorimetric assay [6,7]. Briefly, the test samples were first dissolved in DMSO/MHB. The solution obtained was then added to MHB, and serially diluted two fold (in a 96-wells microplate). One hundred microlitres (100 μ l) of inoculum (1.5×10^6 CFU/ml) prepared in MHB was then added. The plates were covered with a sterile plate sealer, then agitated to mix the contents of the wells using a shaker and incubated at 37°C for 18 h. The final concentration of DMSO was lower than 2.5% and did not affect the microbial growth. Wells containing MHB, 100 μ l of inoculum and DMSO at a final concentration of 2.5% served as negative control (this internal control was systematically added). The total volume in each well was 200 μ l. Chloramphenicol was used as reference antibiotic. The MICs of samples were detected after 18 h incubation at 37°C, following addition (40 μ l) of 0.2 mg/ml INT and incubation at 37°C for 30 minutes. Viable bacteria reduced the yellow dye to pink. MIC was defined as the lowest sample concentration that prevented this change and exhibited complete inhibition of microbial growth [8].

Samples were tested alone and then, in the presence of PA β N at 30 μ g/ml final concentration. Two of the

Table 1 Spices used in the present study and evidence of their activities.

Spice samples (Family)	Herbarium Voucher number ^a	Part used	Bioactive (or potentially active) compounds ^b and screened activity ^c for crude plant extract
<i>Fagara xanthozyloides</i> (Watern. (Rutaceae))	21793/HNC/SRF	Fruits	<u>Antimicrobial activity of essential oil</u> [S: Ec, Bc, Bs, Af, Kp, Sa, Sf [19]; <u>Cytotoxicity of fruits crude methanol extract</u> [weak activity on leukemia CCRF-CEM and CEM/ADR5000 cells, and pancreatic MiaPaCa-2 cell lines] [27]
<i>Dichrostachys glomerata</i> (Forsk) chuov (Mimosaceae)	15220/SRF-Cam	Bark, fruits	<u>Cytotoxicity of roots crude methanol extract</u> [weak activity on leukemia CCRF-CEM and CEM/ADR5000 cells, and pancreatic MiaPaCa-2 cell lines] [27]
<i>Aframomum citratum</i> (Pereira). Schum (Zingiberaceae)	37736/SRF-Cam	Leaves, fruits	<u>Cytotoxicity of leaves crude methanol extract</u> [weak activity on leukemia CCRF-CEM and CEM/ADR5000 cells, and pancreatic MiaPaCa-2 cell lines] [27]
<i>Beilschmiedia cinnamomea</i> (Stapf) Robyns & Wilczek (Lauraceae)	6933/SRF-Cam	Roots	/
<i>Echinops giganteus</i> A. Rich. (Asteraceae)	23647/SRF-Cam	Rhizomes	<u>Antimicrobial</u> [lupeol sitosteryl; β -D-glucopyranoside] [28-31]; <u>Cytotoxicity of rhizome crude methanol extract</u> [Significant activity with IC ₅₀ values of 6.68; 7.96 and 9.84 μ g/ml respectively on leukemia CCRF-CEM cells, CEM/5000 cells and pancreatic MiaPaCa-2 cell lines] [27]
<i>Mondia whitei</i> (Hook F). Skell. (Periplocaceae)	42920/HNC	Fruits	<u>Reproduction system</u> [Roots water extract (400 mg/kg/day) for 55 days caused testicular lesions resulting in the cessation of spermatogenesis, degenerative changes in the somniferous tubules and epididymides in rats] [32]
<i>Olex subscorpioidea</i> Oliv. (Olacaceae)	3528/SRFK	Seeds	<u>Antibacterial and cytotoxic against <i>Artemia salina</i></u> [Santalbic acid] [33,34]; <u>Cytotoxicity of leaves crude methanol extract on cancer cells</u> [weak activity on leukemia CCRF-CEM and pancreatic MiaPaCa-2 cell lines and significant activity with IC ₅₀ of 10.65 μ g/ml on CEM/ADR5000 cells] [27]
<i>Solanum melongena</i> L.Var inerme D.C Hiern. (Solanaceae)	22615/SRFC	Fruits	<u>Antimicrobial activity of methanol, dichloromethane and petrol ether extracts of the fruits</u> : [Q: Tm, Tr, Tt, Ca et Tb [35]
<i>Piper capense</i> Lin.f (Piperaceae)	7650/SRF-Cam	Fruits	<u>Insecticidal</u> [N-isobutyl-11-(3, 4-methylenedioxyphenyl)-2E, 4E, 10E-undecatrienamamide; N-pyrrolidyl-12-(3, 4-methylene-dioxyphenyl)-2E, 4E, 9E, 11Z-dodecatetraenamamide; N-isobutyl-13-(3, 4-methylenedioxyphenyl)-2E, 4E, 12E-tridecatrienamamide; N-isobutyl-2E, 4E-decadienamamide; N-isobutyl-2E, 4E-dodecadienamamide] [36]; <u>Cytotoxicity of fruit crude methanol extract</u> [Significant activity with IC ₅₀ values of 7.02; 6.56 and 8.92 μ g/ml respectively on leukemia CCRF-CEM cells, CEM/5000 cells and pancreatic MiaPaCa-2 cell lines] [27]
<i>Xylopia aethiopica</i> (Dunal) A. Rich. (Annonaceae)	16419/SRF-Cam	Bark, leaves, roots, seeds	<u>Antimicrobial</u> [volatile oil of seeds] [19]; <u>Antioxidant</u> [volatile oil of seeds] [37]; <u>Cytotoxicity of seeds crude methanol extract</u> [Significant activity with IC ₅₀ values of 3.91; 7.4 and 6.86 μ g/ml respectively on leukemia CCRF-CEM cells, CEM/5000 cells and pancreatic MiaPaCa-2 cell lines] [27]
<i>Scorodophloeus zenkeri</i> Harms. (Caesalpinaceae)	44803/SRF-Cam	Bark	<u>Antimicrobial</u> : [2, 4, 5, 7-Tetrathiaoctane; 2, 4, 5, 6, 8-pentathianonane; 2, 3, 4, 6, 8-pentathianonane; 2, 3, 5, 6, 8, 10-hexathiaundecane; 2, 3, 5-trithiahexane 5-oxide; 2, 4, 5, 7-tetrathiaoctane 2-oxide; 2, 3, 5, 7-tetrathiaoctane 3, 3-dioxide; 2, 3, 5-trithiahexane 3, 3-dioxide [38]; <u>Cytotoxicity of bark crude methanol extract on cancer cells</u> [weak activity on leukemia CCRF-CEM and pancreatic MiaPaCa-2 cell lines and significant activity with IC ₅₀ of 10.65 μ g/ml on CEM/ADR5000 cells] [27]

^a(HNC): Cameroon National Herbarium; (SRF): Société des réserves forestières; Cam: Cameroon; ^b(/): Not reported

^c[Screened activity: significant (S: CMI < 100 μ g/ml), moderate (M: 100 < CMI \leq 625 μ g/ml), Weak (W: CMI > 625 μ g/ml) Q: Qualitative activity based on the determination of inhibition zone [11,12]. Tm: *Trichophyton mentagophytes*; Tr: *Trichophyton rubrum*; Tt: *Trichophyton tonsurans* Tb: *Trichosporon beigellii*; Ca: *Candida albicans*; Ck: *Candida krusei*; Af: *Aspergillus flavus*; Bc: *Bacillus cereus*; Bs: *Bacillus subtilis*; Ec: *Escherichia coli*; Kp: *Klebsiella pneumoniae*; Sa: *Staphylococcus aureus*; Sf: *Streptococcus faecalis*.

best extracts [those from *D. glomerata* and *B. cinnamomea*] were also tested in association with antibiotics at MIC/2 and MIC/5. These concentrations were selected following a preliminary assay on one of the tested MDR bacteria, *P. aeruginosa* PA124 (See Additional file 1, Table A1). All assays were performed in triplicate and repeated thrice. Fractional inhibitory concentration (FIC) was calculated as the ratio of MIC_{Antibiotic in combination}/MIC_{Antibiotic alone} and the interpretation made as follows: synergistic (<0.5), indifferent (0.5 to 4), or antagonistic (>4) [9] (The FIC values available in Additional file 1, Tables A2 and A3).

Results

Phytochemical composition of the spice extracts

The results of the phytochemical studies (Table 3) showed that all the tested extracts contain alkaloids, phenols and tannins. Anthocyanins, anthraquinones, flavonoids, saponins, sterols and triterpenes were selectively present.

Antibacterial activity of the spice extracts

The results of the antibacterial activity of the extract alone on a panel of Gram negative bacteria are summarized in Table 4. It appears that the extract from

Table 2 Bacterial strains and features

Strains	Features	References
<i>Escherichia coli</i>		
ATCC8739 and ATCC10536	Reference strains	
AG100	Wild-type <i>E. coli</i> K-12	[39]
AG100A	AG100 Δ acrAB::KAN ^R	[39]
AG100A _{TET}	Δ acrAB mutant de AG100, avec le gène <i>acrF</i> sur-exprimé; TET ^R	[39]
AG102	Δ acrAB mutant AG100, owing <i>acrF</i> gene markedly over-expressed; TET ^R	[40]
MC4100	Wild type <i>E. coli</i>	
W3110	Wild type <i>E. coli</i>	[41]
<i>Enterobacter aerogenes</i>		
ATCC13048	Reference strains	
EA-CM64	CHL ^R resistant variant obtained from ATCC13048 over-expressing the AcrAB pump	[42]
EA3	Clinical MDR isolate; CHL ^R , NOR ^R , OFX ^R , SPX ^R , MOX ^R , CFT ^R , ATM ^R , FEP ^R	[42]
EA27	Clinical MDR isolate exhibiting energy-dependent norfloxacin and chloramphenicol efflux with KAN ^R AMP ^R NAL ^R STR ^R TET ^R	[42,43]
EA289	KAN sensitive derivative of EA27	[43,44]
EA294	EA289 <i>acrA</i> ::KAN ^R	[44]
EA298	EA 289 <i>tolC</i> ::KAN ^R	[44]
<i>Enterobacter cloacae</i>		
ECCI69	Clinical isolates	Laboratory collection of UNR-MD1, University of Marseille, France
BM47	Clinical isolates	Laboratory collection of UNR-MD1, University of Marseille, France
BM67	Clinical isolates	Laboratory collection of UNR-MD1, University of Marseille, France
<i>Klebsiella pneumoniae</i>		
ATCC12296	Reference strains	
KP55	Clinical MDR isolate, TET ^R , AMP ^R , ATM ^R , CEF ^R	[45]
KP63	Clinical MDR isolate, TET ^R , CHL ^R , AMP ^R , ATM ^R	[45]
K24	AcrAB-TolC	Laboratory collection of UNR-MD1, University of Marseille, France
K2	AcrAB-TolC	Laboratory collection of UNR-MD1, University of Marseille, France
<i>Providencia stuartii</i>		
NEA16	Clinical MDR isolate, AcrAB-TolC	
ATCC29914	Clinical MDR isolate, AcrAB-TolC	[46]
PS2636	Clinical MDR isolate, AcrAB-TolC	
PS299645	Clinical MDR isolate, AcrAB-TolC	
<i>Pseudomonas aeruginosa</i>		
PA 01	Reference strains	
PA 124	MDR clinical isolate	[26]

^aAMP, ATM^R, CEF^R, CFT^R, CHL^R, FEP^R, KAN^R, MOX^R, STR^R, TET^R. Resistance to ampicillin, aztreonam, cephalothin, cefadroxil, chloramphenicol, cefepime, kanamycin, moxalactam, streptomycin, and tetracycline; MDR: Multidrug resistant.

D. glomerata was able to prevent the growth of all the twenty nine tested bacteria with MIC \leq 1024 μ g/ml. All other samples showed selective activity; their inhibitory activity being recorded on 28 of the 29 (96.6%) tested bacteria for *B. cinnamomea*, 24/29 (82.8%) for *A. citratum*, 19/29 (62.5%) for *P. capense*, 18/29 (62.1%) for *E.*

giganteus and *F. Xanthoxyloides*, 15/29 (51.7%) for *O. subscorpioidea*, 13/29 (44.8%) for *X. aethiopica*, 12/29 (41.4%) for *M. whitei*, 6/29 (20.7%) for *S. melongena* and 4/29 (13, 79%) for *S. zenkeri*.

MIC values below 100 μ g/ml (Table 4) were recorded with the extract of *B. cinnamomea* against *Enterobacter*

Table 3 Extraction yields, aspects and phytochemical composition of the plant extracts

Spice samples	Extraction yield*	Physical aspect	Phytochemical composition								
			Alkaloids	Anthocyanins	Anthraquinones	Flavonoids	Phenols	Saponins	Tannins	Sterols	Triterpenes
<i>Fagara xanthoxyloides</i>	12.13	Oily brown	+	-	+	+	+	-	+	-	-
<i>Dichrostachys glomerata</i>	18.29	Brown paste	+	+	+	+	+	+	+	+	+
<i>Aframomum citratum</i>	16.32	Brown paste	+	-	-	+	+	+	+	+	-
<i>Beilschmiedia cinnamomea</i>	5.67	Black paste	+	+	+	+	+	-	+	-	+
<i>Echinops giganteus</i>	8.87	Oily brown	+	+	+	+	+	-	+	-	+
<i>Mondia whitei</i>	7.33	Brown paste	+	+	+	+	+	+	+	+	+
<i>Olox subscorpioidea</i>	12.34	Brown paste	+	-	+	+	+	-	+	-	+
<i>Solanum melongena</i>	14.30	Black paste	+	+	+	+	+	+	+	+	+
<i>Piper capense</i>	12.87	Brown paste	+	-	-	-	+	+	+	+	+
<i>Xylopi aethiopica</i>	26.42	Brown paste	+	-	-	-	+	+	+	-	+
<i>Scorodophloeus zenkeri</i>	4.67	Brown paste	+	-	-	-	+	+	+	-	+

(+): Present; (-): Absent; *The yield was calculated as the ratio of the obtained methanol extract according to the initial mass of the spice powder

Table 4 Minimal inhibitory concentration (MIC) of the studied spice extracts and CHL on the studied bacterial species

Bacterial strains	Tested samples and MIC in µg/ml in the absence and presence of PAβN (in parenthesis)											
	<i>F. xanthoxyloides</i>	<i>D. glomerata</i>	<i>A. citratum</i>	<i>B. cinnamomea</i>	<i>E. giganteus</i>	<i>M. whitei</i>	<i>S. melongena</i>	<i>O. subscorpioidea</i>	<i>P. capense</i>	<i>X. aethiopica</i>	<i>S. zenkeri</i>	CHL
<i>E. coli</i>												
ATCC8739	- (1024)	1024 (512)	1024 (256)	- (256)	- (-)	- (1024)	- (128)	- (-)	- (512)	- (512)	- (-)	4 (< 2)
ATCC10536	1024	512 (128)	1024	1024	1024	1024	-	512	256	64	256	< 2 (< 2)
AG100	256	512 (256)	1024 (1024)	1024 (1024)	1024 (1024)	1024 (-)	1024 (-)	512 (-)	- (1024)	1024 (1024)	- (-)	8 (< 2)
AG100A	1024	1024 (256)	1024	1024	1024	-	-	1024	1024	-	-	< 2 (< 2)
AG100A _{TET}	-	1024 (512)	1024	512	1024	-	-	-	1024	1024	-	64 (< 2)
AG102	1024	512 (256)	1024	512	1024	-	-	1024	1024	1024	-	32 (< 2)
MC4100	256	256(< 8)	512	256	512	1024	1024	1024	1024	1024	-	32
W3110	- (512)	256 (< 8)	512 (< 8)	1024 (< 8)	1024 (1024)	1024 (512)	-	512 (512)	1024 (1024)	- (-)	- (-)	4 (< 2)
<i>E. aerogenes</i>												
ATCC13048	-	512 (512)	1024	512	-	-	-	-	-	-	-	8 (< 2)
CM64	- (-)	512 (512)	- (1024)	1024 (1024)	1024 (1024)	1024 (-)	- (-)	- (-)	- (-)	- (-)	- (-)	256 (8)
EA3	1024	1024 (1024)	-	1024	-	-	-	1024	1024	-	-	- (128)
EA27	1024	512 (256)	1024	1024	1024	1024	-	512	1024	1024	-	256 (< 2)
EA289	-	1024 (1024)	1024	512	1024	-	-	-	-	-	-	- (64)
EA298	-	1024 (128)	-	1024	-	1024	-	256	512	1024	128	64 (< 2)
EA294	256	128	512	64	256	-	1024	256	256	256	512	8
<i>E. cloacae</i>												
ECC169	512 (-)	1024 (1024)	1024 (1024)	1024 (512)	1024 (-)	- (-)	- (-)	- (-)	-	-(1024)	- (256)	- (16)
BM47	1024 (1024)	1024 (256)	1024 (1024)	1024 (1024)	1024 (-)	- (-)	- (-)	- (-)	512(64)	1024(1024)	- (-)	- (< 2)
BM67	1024 (1024)	1024 (128)	1024	1024	-	-	-	-	512	1024	-	256 (16)
<i>K. pneumoniae</i>												
ATCC11296	1024 (1024)	512 (128)	512 (512)	256 (256)	1024 (256)	- (1024)	- (1024)	1024 (512)	1024 (256)	- (-)	- (-)	4 (< 2)
KP55	1024	512 (256)	1024	512	-	-	-	1024	1024	-	-	64 (2)
KP63	1024	512 (< 8)	256	512	512	1024	512	256	256	64	-	64 (< 2)
K24	512	512 (32)	512	256	32	1024	1024	-	1024	-	-	16 (< 2)
K2	-	1024 (128)	-	1024	-	-	-	1024	512	-	-	32 (4)
<i>P. stuartii</i>												
NEA16	1024	512 (32)	128	256	1024	1024	1024	512	256	512	1024	32 (8)
ATCC29914	1024	1024 (512)	512	512	1024	1024	-	-	1024	-	-	16 (8)
PS2636	-	512	1024	1024	-	-	-	-	-	-	-	32
PS299645	1024 (-)	256 (256)	256	256 (512)	- (-)	- (-)	- (512)	- (-)	- (-)	- (-)	- (-)	32 (< 2)
<i>P. aeruginosa</i>												
PA01	-	1024 (512)	1024	1024	-	-	-	-	-	1024	-	16 (< 2)
PA124	-	512 (512)	-	1024	-	1024	-	-	-	-	-	32 (< 2)

(-): MIC not detected at up to 1024 µg/ml for the les extracts and 256 µg/ml for CHL. (): values in parenthesis are MIC of substance in the presence of PAβN at 30 µg/ml. The MIC of PAβN was 64 µg/ml on *E. coli*, AG100A, 512 µg/ml on ATCC11296, BM67, EA27, EA289; 1024 µg/ml on AG100A_{TET}, ATCC13048, CM64; and > 1024 µg/ml on other bacteria. CHL: chloramphénicol

aerogenes EA294 (64 µg/ml), *E. giganteus* on *Klebsiella pneumoniae* K24 (32 µg/ml) and *X. aethiopica* on *Escherichia coli* ATCC10536 and *Klebsiella pneumoniae* KP63 (64 µg/ml).

Role of efflux pumps in susceptibility of Gram negative bacteria to the tested spice extracts

The various strains and MDR isolates were tested for their susceptibilities to the spice extracts, and reference antibiotic, CHL in the presence of PAβN, a well-known efflux pump inhibitor. The results presented in Table 4 showed that the activity of the extract from *D. glomerata* significantly increased in the presence of PAβN on 18/26 (69.2%) of the tested bacteria. The MIC values below 100 µg/ml were noted with this extract against *E. coli* MC4100 and W3110 (< 8 µg/ml), *K. pneumoniae* KP63 and K24 (< 8 µg/ml and 32 µg/ml respectively) and *P. stuartii* NAE16 (32 µg/ml). Apart from the extract of *D. glomerata*, PAβN did not induce an increased activity of other tested extract.

Effects of the association of some spice extracts with antibiotics

To evaluate the possible synergistic effects of the extracts with antibiotics, four of the most active samples (*F. xanthoxyloides*, *D. glomerata*, *B. cinnamomea* and *O. subscorpioidea*) were selected. A preliminary study using *P. aeruginosa* PA124, one of the MDR bacteria used in this work, was carried out with ten antibiotics (CLX, AMP, ERY, KAN, CHL, TET, FEP, STR, CIP and NOR) to select the appropriate sub-inhibitory concentrations to be used. The results (see Additional file 1, Table A1) allow the selection of MIC/2 and MIC/5 as the sub-inhibitory concentrations of the extracts from *D. glomerata* and *B. cinnamomea*, which were then tested on eight MDR bacteria, *E. coli* AG100, AG100_{TET}, *K. pneumoniae* KP55, *E. aerogenes* EA3, EA27, EA289, CM64 in addition to *P. aeruginosa* PA124. The results are summarized in Tables 5 and 6. Synergistic effects were observed with the association between *D. glomerata* (Table 5, Additional file 1, Table A2) and *B. cinnamomea* (Table 6, Additional file 1, Table A3) and most of the antibiotics on the studied MDR bacteria. At MIC/2, synergistic effects were noted with the extract of *D. glomerata* on 25% (2/8) of the tested bacteria for CLX and AMP, 50% (4/8) for KAN, 62.5% (5/8) for CHL, FEP, STR, CIP, 75% (6/8) for ERY and 87.5% (7/8) for NOR and TET. Increase in MIC values of > 8 fold were recorded at MIC/2 with CHL, TET, STR, CIP, NOR (Table 5). At MIC/5, synergistic effects were noted on 50% of the eight tested MDR bacteria in the case of STR and CIP, 62.5% in the case of ERY and 75% in the case of CHL, TET and NOR.

The extract of *B. cinnamomea* at MIC/2 (Table 6) also induced significant increase of the activity of several

antibiotics, the synergistic effects being noted on 25% of the tested bacteria in the case of CLX and AMP, 50% in the case of KAN, 62.5% in the case of FEP and STR, 75% in the case of CHL, TET and CIP, 87.5% in the case of ERY and 100% for NOR. With this extract, synergistic effects were also observed at MIC/5 on 25% of the studied MDR bacteria in the case of CLX and AMP, 37.5% in the case of STR and KAN, 50% in the case of TET, 62.5% in the case of CHL, FEP, NOR and CIP and 75% in the case of ERY.

Discussion

Phytochemical composition of the spice extracts

Phytochemical screening revealed the presence of several classes of secondary metabolites. Though the detection of such metabolites does not automatically predict the antimicrobial activity of a plant extract, it has clearly been demonstrated that several compounds belonging to the investigated classes of metabolites showed antibacterial activities [4,10-12].

Antibacterial activity of the spice extract

Phytochemicals are routinely classified as antimicrobials on the basis of susceptibility tests that produce MIC in the range of 100 to 1000 µg/ml [13]. Activity is considered to be significant if MIC values are below 100 µg/ml for crude extract and moderate when 100 < MIC < 625 µg/ml [11]. Therefore, the activity recorded with *B. cinnamomea* and *E. giganteus* respectively on *E. aerogenes* EA294 and *K. pneumoniae* K24, and *X. aethiopica* on *E. coli* ATCC10536 and *K. pneumoniae* KP63 can be considered significant. Alternative criteria have been described by Fabry et al. [14], which consider extracts having MIC values below 8000 µg/ml to have noteworthy antimicrobial activity. Under these less stringent criteria, and considering the fact that the spices tested are used as food ingredients with limited toxicity, the overall activity recorded with several extracts, most notably those of *D. glomerata*, *B. cinnamomea*, *A. citratum*, *P. capense*, *E. giganteus*, *F. Xanthoxyloides* and *O. subscorpioidea*, could be considered important. Besides, some of the tested samples were more active than CHL used as reference antibiotic on some of the MDR bacteria such as *E. cloacae* ECC169 and BM47, *E. aerogenes* EA27 and EA289, highlighting the importance of the results reported herein. It can be noted that all the investigated phytochemical classes were detected in the extracts of *D. glomerata*, *S. melongena* and *M. withei*. Contrary to *D. glomerata* extract that exhibited a good spectrum of activity, the inhibition potential of *S. melongena* and *M. withei* was lower and seems not in correlation with their chemical composition. This clearly confirms the fact that the presence of secondary metabolites does not automatically predict the antimicrobial activity of a plant

Table 5 Minimal inhibitory concentration (MIC) in µg/ml of antibiotics in the absence and presence of the sub-inhibitory concentrations of *D. glomerata* extracts against MDR bacteria.

Bacterial strains	Antibiotics and MIC in absence and presence <i>D. glomerata</i> extracts at MIC/2 and MIC/5														
	Chloramphenicol			Cloxacillin			Ampicillin			Erythromycin			Kanamycin		
	Alone	MIC/2	MIC/5	Alone	MIC/2	MIC/5	Alone	MIC/2	MIC/5	Alone	MIC/2	MIC/5	Alone	MIC/2	MIC/5
PA124	32	32 (1) ^I	16 (2) ^S	-	-	-	64	-	-	64	32 (2) ^S	64 (1) ^I	64	16 (4) ^S	64 (1)
CM64	256	256 (1) ^I	-	-	-	-	-	-	256	-	64	128	1	1 (1) ^I	1 (1) ^I
EA3	-	32 (> 16) ^S	32 (> 16) ^S	-	256	256	-	-	-	64	32 (2) ^S	64 (1) ^I	32	16 (2) ^S	16 (2) ^S
EA27	256	32 (8) ^S	64 (4) ^S	256	64 (4) ^S	64 (4) ^S	64	64 (1) ^I	64 (1) ^I	32	32 (1) ^I	32 (1) ^I	16	16 (1) ^I	16 (1) ^I
EA289	-	-	-	-	-	-	-	256	64	256	64 (4) ^S	128 (2) ^S	4	< 2 (> 2) ^S	4 (1) ^I
KP55	64	8 (8) ^S	16 (4) ^S	-	-	-	-	-	-	256	128 (2) ^S	128 (2) ^S	32	32 (1) ^I	32 (1) ^I
AG100 _{TET}	64	8 (8) ^S	16 (4) ^S	-	-	-	-	16	32	32	32 (1) ^I	16 (2) ^S	32	2 (16) ^S	8 (4) ^S
AG100	8	< 2 (> 4) ^S	< 2 (> 4) ^S	256	128 (2) ^S	64 (4) ^S	64	4 (16) ^S	4 (16) ^S	32	< 2 (> 16) ^S	4 (8) ^S	< 2	< 2	< 2
Bacterial strains	Tetracycline			Cefepime			Streptomycin			Ciprofloxacin			Norfloxacin		
	Alone	MIC/2	MIC/5	Alone	MIC/2	MIC/5	Alone	MIC/2	MIC/5	Alone	MIC/2	MIC/5	Alone	MIC/2	MIC/5
PA124	4	< 0, 5 (> 8) ^S	2 (2) ^S	-	-	-	16	16 (1) ^I	16 (1) ^I	16	< 0, 5 (> 8) ^S	16 (1) ^I	128	-	-
CM64	32	4 (8) ^S	8 (4) ^S	256	64 (4) ^S	128 (2) ^S	8	< 2 (> 4) ^S	4 (2) ^S	1	1 (1) ^I	1 (1) ^I	2	1 (2) ^S	1 (2) ^S
EA3	2	1 (2) ^S	2 (2) ^S	-	-	-	16	8 (2) ^S	8 (2) ^S	64	4 (16) ^S	64 (1) ^I	64	32 (2) ^S	64 (1) ^I
EA27	16	4 (4) ^S	8 (2) ^S	256	128 (2) ^S	128 (2) ^S	8	4 (2) ^S	4 (2) ^S	2	2 (1) ^I	2 (1) ^I	16	2 (8) ^S	4 (4) ^S
EA289	8	1 (8) ^S	2 (4) ^S	-	256 (> 2) ^S	-	64	8 (8) ^S	32 (2) ^S	32	16 (2) ^S	16 (2) ^S	64	16 (4) ^S	32 (2) ^S
KP55	4	2 (2) ^S	2 (2) ^S	-	128 (> 4) ^S	-	8	8 (1) ^I	8 (1) ^I	128	4 (32) ^S	32 (4) ^S	128	32 (4) ^S	32 (4) ^S
AG100 _{TET}	4	1 (4) ^S	2 (2) ^S	-	32 (> 16) ^S	-	16	2 (8) ^S	16 (1) ^I	64	32 (2) ^S	16 (4) ^S	64	8 (8) ^S	16 (4) ^S
AG100	< 2	< 2 (> 4) ^S	< 2 (> 4) ^S	256	< 2 (> 128) ^S	< 2 (> 128) ^S	256	256 (1) ^I	256 (1) ^I	< 2	< 2	< 2	32	4 (8) ^S	32 (1) ^I

MIC/2: concentration of plant extract added equal to 256 µg/ml for PA124, CM64, EA3, EA27, KP55, AG100; and to 512 µg/ml for EA289 and AG100_{TET}.

MIC/5: concentration of plant extract added equal to 102.4 µg/ml for PA124, CM64, EA3, EA27, KP55, AG100; and to 204.8 µg/ml for EA289 and AG100_{TET}.

(I): Values in bracket are folds increase of activity. S: synergy, I: indifference; (-): > 512 µg/ml

Table 6 Minimal inhibitory concentration (MIC) of antibiotics in the absence and presence of the sub-inhibitory concentrations of *B. cinnamomea* extract ($\mu\text{g/ml}$) against some MDR bacteria.

Bacterial strains	Antibiotics and MIC in absence and presence <i>B. cinnamomea</i> extracts at MIC/2 and MIC/5														
	Chloramphenicol			Cloxacillin			Ampicillin			Erythromycin			Kanamycin		
	Alone	MIC/2	MIC/5	Alone	MIC/2	MIC/5	Alone	MIC/2	MIC/5	Alone	MIC/2	MIC/5	Alone	MIC/2	MIC/5
PA124	32	< 0, 5 (> 32) ^S	32 (1) ^S	-	-	-	-	-	-	64	8 (8) ^S	32 (2) ^S	32	32 (1)	64
CM64	256	-	-	-	-	-	-	-	-	-	64 (> 8) ^S	64 (> 8) ^S	1	< 0.5 (> 2) ^S	< 0.5 (> 2) ^S
EA3	-	16 (> 32) ^S	32	-	256	-	-	-	-	64	8 (8) ^S	32 (2) ^S	32	16 (2) ^S	16 (2) ^S
EA27	256	8 (32) ^S	16 (16) ^S	256	< 0, 5 (> 512) ^S	16(16) ^S	64	64 (1) ^I	64 (1) ^I	32	8 (4) ^S	32 (2) ^S	16	< 0.5 (> 32) ^S	16(1) ^I
EA289	-	256	-	-	-	-	-	-	-	256	32 (8) ^S	64 (4) ^S	4	4 (1) ^I	4 (1) ^I
KP55	64	8 (8) ^S	8 (8) ^S	-	-	-	-	-	-	256	128 (2) ^S	128 (2) ^S	32	32 (1) ^I	32 (1) ^I
AG100 _{TET}	64	16 (4) ^S	32 (2) ^S	-	128 (> 4) ^S	128 (> 4) ^S	-	8 (> 64) ^S	32 (16) ^S	32	32 (1) ^I	32 (1) ^I	32	1 (32) ^S	8 (4) ^S
AG100	8	< 2 (> 4) ^S	4 (2) ^S	256	256 (1)	256 (1)	64	< 2 (> 32) ^S	4 (16) ^S	32	< 2 (> 16) ^S	4 (8) ^S	< 2	< 2	< 2
Bacterial strains	Tetracycline			Cefepime			Streptomycin			Ciprofloxacin			Norfloxacin		
	Alone	MIC/2	MIC/5	Alone	MIC/2	MIC/5	Alone	MIC/2	MIC/5	Alone	MIC/2	MIC/5	Alone	MIC/2	MIC/5
PA124	4	2 (2) ^S	4 (1) ^S	-	64 (> 8) ^S	-	16	16 (1)	16 (1)	16	16 (1) ^I	16 (1) ^I	128	64 (2) ^S	64 (2) ^S
CM64	32	8 (4) ^S	8 (4) ^S	256	32 (8) ^S	32 (8) ^S	8	< 2 (> 4) ^S	< 2 (> 4) ^S	1	1 (1) ^I	1 (1) ^I	2	1 (2) ^S	2 (1) ^I
EA3	2	1 (2) ^S	2 (1) ^I	-	-	-	16	4 (4) ^S	4 (4) ^S	64	4 (16) ^S	8(8) ^S	64	16 (4) ^S	32 (2) ^S
EA27	16	1 (16) ^S	4 (4) ^S	256	16 (16) ^S	64(4) ^S	8	< 0.5 (> 16) ^S	2 (4) ^S	2	< 0.5 (> 4) ^S	2 (1) ^I	16	< 0.5 (> 32) ^S	4 (4) ^S
EA289	8	4 (2) ^S	4 (2) ^S	-	256 (> 2) ^S	256 (> 2) ^S	64	16 (4) ^S	64 (1) ^I	32	16 (2) ^S	16 (2) ^S	64	32 (2) ^S	32 (2) ^S
KP55	4	4 (1) ^I	4 (1) ^I	-	64 (> 8) ^S	128 (> 2) ^S	8	8 (1) ^I	8 (1) ^I	128	8 (16) ^S	16 (8) ^S	128	16 (8) ^S	32 (4) ^S
AG100 _{TET}	4	2 (2) ^S	2 (2) ^S	-	256 (> 2) ^S	64(> 4) ^S	16	8 (2) ^S	16 (1) ^I	64	4 (16) ^S	8 (8) ^S	64	32 (2) ^S	64 (1) ^I
AG100	< 2	< 2	< 2	256	64 (4) ^S	64 (4) ^S	256	256 (1) ^I	256 (1) ^I	4	< 2 (> 2) ^S	< 2 (> 2) ^S	32	4 (8) ^S	32 (1) ^I

MIC/2: concentration of plant extract added equal to 512 $\mu\text{g/ml}$ for PA124, CM64, EA3, EA27, AG100; and to 256 $\mu\text{g/ml}$ for EA289, KP55 and AG100_{TET}
 MIC/5: concentration of plant extract added equal to 204.8 $\mu\text{g/ml}$ for PA124, CM64, EA3, EA27, KP55, AG100; and to 102.4 $\mu\text{g/ml}$ for EA289 and AG100_{TET}
 (I): Values in bracket are folds increase of activity. S: synergy, I: indifference; (-): > 512 $\mu\text{g/ml}$

extract though it is a good indication of its possible pharmacological potential.

To the best of our knowledge, the antibacterial activity of *B. cinnamomea* and *P. capense* is being reported for the first time. Moreover, the present work reports for the first time the activity of the tested spices on MDR bacteria. Nevertheless, the antimicrobial potential of some of the plants or related genus were demonstrated on sensitive strains. Banso and Adeyemo [15] reported the presence of antibacterial tannins in the genus *Dichrostachys*. Chouna et al. [16] also demonstrated that *Beilschmiedia anacardioides* was significantly active against *Bacillus subtilis*, *Micrococcus luteus* and *Streptococcus faecalis*. Plants of the genus *Echinops* such as *E. ellenbeckii* and *E. longisetus* were found active on *Staphylococcus aureus* [17] meanwhile the antibacterial activity of the essential oils and alkaloids from *F. xanthoxyloides* was also documented [18,19]. The aqueous and ethanol extracts from *O. subscorpioidea* were found active on both bacteria and fungi [20]. The results obtained in the present work therefore provide additional information on the studied plants and are in consistence with some of the previous reports.

Role of efflux pumps in susceptibility of Gram negative bacteria to the tested spice extracts

Tripartite efflux systems, mainly those clinically described as AcrAB-TolC in *Enterobacteriaceae* or MexAB-OprM in *P. aeruginosa*, are associated with a major human health problem as they play a central role in multidrug resistance of pathogenic Gram negative bacteria [21-23]. PA β N has been reported as a potent inhibitor of the RND efflux systems and is especially active on AcrAB-TolC and MexAB-OprM [22,24,25]. To determine the role of efflux pumps in this work, the concentration of PA β N used (30 μ g/ml) had no intrinsic effect on the bacteria as previously determined [26]. In contrast, with these conditions significant increase of the antibacterial activity of *D. glomerata* extract was noted, showing that one or more active compounds from this plant could be substrate(s) of efflux pumps acting in resistant strains of *E. coli*, *K. pneumoniae* and *P. stuartii*. These data suggest that possible association of the extract of *D. glomerata* and efflux pump inhibitor can be envisaged to improve the fight against MDR phenotypes.

Effects of the association of extracts from *D. glomerata* and *B. cinnamomea* with antibiotics

The association of natural products such as plant extracts and antibiotics constitutes an alternative in the fight against MDR bacteria. Significant synergistic effects were noted with both *D. glomerata* and *B. cinnamomea* extracts when they were associated with several antibiotics. Such effects might be due either to the action of the

active compounds or possible inhibition of the efflux pumps by other compounds of the extracts. The lowest synergistic effects were observed with β -lactamines (CLX and AMP), obviously due to the fact their target are localized in the bacterial cell coat. However, the synergistic effects observed indicate that active compounds of the extract could also present different mode(s) of action from those of the studied antibiotics.

Conclusion

The overall results of the present work provide baseline information for the possible use of the studied spice extracts in the treatment of bacterial infections involving MDR phenotypes. In addition to these antibacterial activities, the data reported herein indicated that possible combinations of the extract of *D. glomerata* with an efflux pump inhibitor, and also the association of extract of this plant as well as that from *B. cinnamomea* with several antibiotics could be used in the control of bacterial infections involving MDR phenotypes.

Additional material

Additional file 1: Table S1. Activities of antibiotics in combination with the sub-inhibitory concentrations of some plants extracts on *Pseudomonas aeruginosa* PA124. **Table S 2.** Fractional Inhibitory Concentrations (FIC) of the association between antibiotics and extract of *D. glomerata* at MIC/2 and MIC/5 (μ g/ml) against MDR bacteria. **Table S 3.** Fractional Inhibitory Concentrations (FIC) of association between antibiotics and extract of *B. cinnamomea* at MIC/2 and MIC/5 (μ g/ml).

Acknowledgements

Authors are thankful to the Cameroon National Herbarium (Yaounde) for plants identification and Mr. Paul K. Lunga and Dr Gerald Ngo Teke for language editing.

Author details

¹Department of Biochemistry, Faculty of science, University of Dschang, Cameroon. ²Transporteurs Membranaires, Chimiorésistance et Drug Design, UMR-MD1, IFR 88, UFRs de Médecine et de Pharmacie, Marseille, France.

Authors' contributions

PAF carried out the study; VK designed the experiments and wrote the manuscript; VK, IKV, JRK and JMP supervised the work; VK and JMP provided the bacterial strains; All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 4 August 2011 Accepted: 1 November 2011

Published: 1 November 2011

References

1. Kamicker BJ, Sweeney MT, Kaczmarek F, Dib-Hajj F, Shang W, Crimin K, Duignan J, Gootz TD: **Bacteria efflux pump inhibitors.** *Methods Mol Med* 2008, **142**:187-204.
2. Lutz JK, Lee J: **Prevalence and Antimicrobial-Resistance of *Pseudomonas aeruginosa* in Swimming Pools and Hot Tubs.** *Int J Environ Res Public Health* 2011, **8**:554-564.
3. Sofowora A: **1993. Medicinal Plants and Traditional Medicine in Africa.** Ibadan, Spectrum Books Limited; 1993.

4. Cowan MM: Plant products as antimicrobial agents. *Clinical Microbiology Reviews* 1999, **12**:564-582.
5. Harborne JB: *Phytochemical Methods*. New York, Chapman and Hall; 1973.
6. Eloff JN: A sensitive and quick microplate method to determine the minimal inhibitory concentration of plant extracts for bacteria. *Planta Med* 1998, **64**:711-713.
7. Mativandele SPN, Lall N, Meyer JJM: Antifungal and antitubercular activity of (the roots of) *Pelargonium reniforme* (CURT) and *Pelargonium sidoides* (DC) (Geraniaceae) root. *S Afr J Bot* 2006, **72**:232-237.
8. Kuete V, Ngameni B, Fotso Simo CC, Kengap Tankeu R, Tchaleu Ngadjui B, Meyer JJM, Lall N, Kuate JR: Antimicrobial activity of the crude extracts and compounds from *Ficus chlamydorcarpa* and *Ficus cordata* (Moraceae). *J Ethnopharmacol* 2008, **120**:17-24.
9. Coutinho HD, Vasconcellos A, Freire-Pessôa HL, Gadelha CA, Gadelha TS, Almeida-Filho GG: Natural products from the termite *Nasutitermes corniger* lower aminoglycoside minimum inhibitory concentrations. *Pharmacogn Mag* 2010, **6**:1-4.
10. Bruneton J: Pharmacognosie: Phytochimie, Plantes medicinales. Paris, Tec & Doc; 3 1999, 263-309.
11. Kuete V: Potential of Cameroonian plants and derived-products against microbial infections: A review. *Planta Med* 2010, **76**:1479-1491.
12. Kuete V, Efferth T: Cameroonian medicinal plants: Pharmacology and derived natural products. *Front Pharmacol* 2010, **1**:1-19.
13. Simões M, Bennett RN, Rosa EA: Understanding antimicrobial activities of phytochemicals against multidrug resistant bacteria and biofilms. *Nat Prod Rep* 2009, **26**:746-757.
14. Fabry W, Okemo PO, Ansorg R: Antibacterial activity of East African medicinal plants. *J Ethnopharmacol* 1998, **60**:79-84.
15. Banso A, Adeyemo SO: Evaluation of antibacterial properties of tannins isolated from *Dichrostachys cinerea*. *Afr J Biotech* 2007, **6**:1785-1787.
16. Chouna JR, Nkeng-Efouet PA, Lenta BN, Devkota PK, Neumann B, Stammeler HG, Kimbu SF, Sewald N: Antibacterial endiandric acid derivatives from *Beilschmiedia anacardioides*. *Phytochemistry* 2009, **70**:684-688.
17. Hymete A, Iversena TH, Rohloff J, Erkob B: Screening of *Echinops ellenbeckii* and *Echinops longisetus* for biological activities and chemical constituents. *Phytomedicine* 2005, **12**:675-679.
18. Chaaib F, Queiroz EF, Ndjoko K, Diallo D, Hostettmann K: Antifungal and antioxidant compounds from the root bark of *Fagara xanthoxyloides*. *Planta Med* 2003, **64**:616-620.
19. Tatsadjieu LN, Essia Ngang JJ, Ngassoum MB, Etoa FX: Antibacterial and antifungal activity of *Xylopiya aethiopicum*, *Monodora myristica*, *Zanthoxylum xanthoxyloides* and *Zanthoxylum lepreurii* from Cameroon. *Fitoterapia* 2003, **74**:469-472.
20. Ayandele AA, Adebisi AO: The phytochemical analysis and antimicrobial screening of extracts of *Oxalis subsoparioides*. *Afr J Biotech* 2007, **6**:868-870.
21. Blot S, Depuydt P, Vandewoude K, De Bacquer D: Measuring the impact of multidrug resistance in nosocomial infection. *Curr Opin Infect Dis* 2007, **20**:391-396.
22. Pietras A, Bavro VN, Furnham N, Pellegrini-Calace M, Milner-White EJ, Luisi BF: Structure and mechanism of drug efflux machinery in Gram negative bacteria. *Curr Drug Target* 2008, **9**:719-728.
23. Papadopoulos CJ, Carson CF, Chang BJ, Riley TV: Role of the MexAB-OprM efflux pump of *Pseudomonas aeruginosa* in tolerance to tea tree (*Melaleuca alternifolia*) oil and its monoterpene components terpinen-4-ol, 1, 8-cineole and α -terpineol. *Appl Environ Microbiol* 2008, **74**:1932-1935.
24. Lomovskaya O, Bostian KA: Practical applications and feasibility of efflux pump inhibitors in the clinic—a vision for applied use. *Biochem Pharmacol* 2006, **71**:910-918.
25. Pagès J-M, Lavigne JP, Leflon-Guibout V, Marcon E, Bert F, Noussair L, Nicolas-Chanoine MH: Efflux Pump, the Masked Side of β -Lactam Resistance in *Klebsiella pneumoniae* Clinical Isolates. *PLoS ONE* 2009, **4**: e4817.
26. Lorenzi V, Muselli A, Bernardini AF, Berti L, Pagès JM, Amaral L, Bolla JM: Geraniol restores antibiotic activities against multidrug-resistant isolate from Gram-negative species. *Antimicrob Agents Chemother* 2009, **53**:2209-2211.
27. Kuete V, Krusche B, Youns M, Voukeng I, Fankam AG, Tankeo S, Lacmata S, Efferth T: Cytotoxicity of some Cameroonian spices and selected medicinal plant extracts. *J Ethnopharmacol* 2011, **134**:803-812.
28. Kojima H, Sato N, Hatano A, Ogura H: Sterol glucosides from *Prunella vulgaris*. *Phytochemistry* 1990, **29**:2351-2355.
29. Tane P, Bergquist K-E, Tene M, Ngadjui BT, Ayafor JF, Sterner O: Cyclodione, an unsymmetrical dimeric diterpene from *Cylicodiscus gabunensis*. *Tetrahedron* 1995, **51**:11595-11600.
30. Kuete V, Eyang KO, Folefoc GN, Beng VP, Hussain H, Krohn K, Nkengfack AE: Antimicrobial activity of the methanolic extract and of the chemical constituents isolated from *Newbouldia laevis*. *Pharmazie* 2007, **62**:552-556.
31. Kuete V, Wansi JD, Mbaveng AT, Kana Sop MM, Tcho Tadjong A, Beng VP, Etoa FX, Wandji J, Marion Meyer JJ, Lall N: Antimicrobial activity of the methanolic extract and compounds from *Teclea afzelii* (Rutaceae). *S Afr J Bot* 2008, **74**:572-576.
32. Watcho P, Kamtchouing P, Sokeng S, Moundipa PF, Tantchou J, Essame JL, Koueta N: Reversible antispermatogenic and antifertility activities of *Mondia whitei* L. in male albino rat. *Phytother Res* 2001, **15**:26-29.
33. Jones GP, Rao KS, Tucker DJ, Richardson B, Barnes A, Rivett DE: Antimicrobial activity of santalbic acid from the oil of *Santalum acuminatum* (Quandong). *Int J Pharmacogn* 1995, **33**:120-123.
34. Cantrell CL, Berhow MA, Phillips BS, Duval SM, Weisleder D, Vaughn SF: Bioactive crude plant seed extracts from the NCAUR oilseed repository. *Phytomedicine* 2003, **10**:325-333.
35. Das J, Lahan JP, Srivastava RB: *Solanum melongena*: A potential source of antifungal agent. *Indian J Microbiol* 2008, **50**:62-69.
36. Gbewonyo WSK, Candy DJ: Chromatographic isolation of insecticidal amides from *Piper guineense* roots. *J Chromatogr A* 1992, **607**:105-111.
37. N'dri K, Bosson KA, Mamyrbekova-Bekro JA, Jean N, Bekro YA: Chemical composition and antioxidant activities of essential. *Eur J Sci Res* 2009, **37**:311-318.
38. Kouokam JC, Jahns T, Becker H: Antimicrobial activity of the essential oil and some isolated sulfur-rich compounds from *Scorodophloeus zenkeri*. *Planta Med* 2002, **68**:1082-1087.
39. Viveiros M, Jesus A, Brito M, Leandro C, Martins M, Ordway D, Molnar M, Molnar J, Amaral L: Inducement and Reversal of Tetracycline Resistance in *Escherichia coli* K-12 and Expression of Proton Gradient-Dependent Multidrug Efflux Pump Genes. *Antimicrob Agents Chemother* 2005, **49**:3578-3582.
40. Elkins CA, Mullis LB: Substrate competition studies using whole-cell with the major tripartite multidrug efflux pumps of *Escherichia coli*. *Antimicrob Agents Chemother* 2007, **51**:923-929.
41. Baglioni P, Bini L, Liberatori S, Pallini V, Marri L: Proteome analysis of *Escherichia coli* W3110 expressing an heterologous sigma factor. *Proteomics* 2003, **3**:1060-1065.
42. Ghisalberti D, Masi M, Pagès J-M, Chevalier J: Chloramphenicol and expression of multidrug efflux pump in *Enterobacter aerogenes*. *Biochem Biophys Res Commun* 2005, **328**:1113-1118.
43. Malla M, Mahamoud A, Chevalier J, Alibert-Franco S, Brouant P, Barbe J, Pagès JM: Alkylaminoquinolines inhibit the bacterial antibiotic efflux pump in multidrug-resistant clinical isolates. *Biochem J* 2003, **376**:801-805.
44. Pradel E, Pagès J-M: The AcrAB-ToIC efflux pump contributes to multidrug resistance in the nosocomial pathogen *Enterobacter aerogenes*. *Antimicrob Agents Chemother* 2002, **46**:2640-2644.
45. Chevalier J, Pagès J-M, Eyraud A, Malla M: Membrane permeability modifications are involved in antibiotic resistance in *Klebsiella pneumoniae*. *Biochem Biophys Res Commun* 2000, **274**:496-499.
46. Tran QT, Mahendra KR, Hajjar A, Ceccarelli M, Davin-Regli A, Winterhalter M, Weingart H, Pagès JM: Implication of Porins in β -Lactam Resistance of *Providencia stuartii*. *J Biol Chem* 2000, **285**:32273-32281.

Pre-publication history

The pre-publication history for this paper can be accessed here:
<http://www.biomedcentral.com/1472-6882/11/104/prepub>

doi:10.1186/1472-6882-11-104

Cite this article as: Fankam et al.: Antibacterial activities of selected Cameroonian spices and their synergistic effects with antibiotics against multidrug-resistant phenotypes. *BMC Complementary and Alternative Medicine* 2011 **11**:104.