

RESEARCH ARTICLE

Open Access



The ethyl acetate fraction of corn silk exhibits dual antioxidant and anti-glycation activities and protects insulin-secreting cells from glucotoxicity

Chia-Chuan Chang^{1†}, Wei Yuan^{2†}, Hsiao-Yuh Roan³, Jia-Ling Chang⁴, Hsiu-Chen Huang⁵, Yu-Ching Lee^{6,7}, Huey Jen Tsay⁸ and Hui-Kang Liu^{4,9*}

Abstract

Background: In this study, we aimed to develop a *Stigmata Maydis* (corn silk) fraction with dual bio-activities against oxidative stress and protein glycation to protect β -cells from diabetes-induced failure.

Methods: Corn silk fractions were prepared by partition and chemically characterised by thin-layer chromatography. Free radical scavenging assay, glycation assay, and cell-based viability test (neutral red) were employed to decide the best fraction. Cell death analysis was executed by annexin V/ Propidium iodide staining. Cell proliferation was measured by WST-1. Finally, β -cell function was evaluated by β -cell marker gene expression (RT-PCR) and acute insulin secretion test.

Results: Four corn silk fractions were prepared from an ethanolic crude extract of corn silk. In vitro assays indicate ethyl acetate fraction (YMS-EA) was the most potent fraction. YMS-EA also attenuated the hydrogen peroxide- or methylglyoxal-induced induction of reactive oxygen species, reduction of cell viability, and inhibition of cell proliferation. However, YMS-EA was unable to prevent hydrogen peroxide-induced apoptosis or advanced glycation end-products-induced toxicity. Under hyperglycemic conditions, YMS-EA effectively reduced ROS levels, improved mRNA expression of insulin, glucokinase, and PDX-1, and enhanced glucose-stimulated insulin secretion. The similarity of bioactivities among apigenin, luteolin, and YMS-EA indicated that dual activities of YMS-EA might be derived from those compounds.

Conclusions: We concluded that YMS-EA fraction could be developed as a preventive food agent against the glucotoxicity to β -cells in Type 2 diabetes.

Keywords: *Stigmata Maydis* (corn silk), Glucotoxicity, Methylglyoxal, Advanced glycation end products, Reactive oxygen species, β -cell failure

* Correspondence: hk.liu@nricm.edu.tw

[†]Equal contributors

⁴Division of Basic Chinese Medicine, National Research Institute of Chinese Medicine, Ministry of Health and Welfare, Taipei, Taiwan, ROC

⁹Ph.D Program for the Clinical Drug Discovery from Herbal Medicine, College of Pharmacy, Taipei Medical University, Taipei, Taiwan, ROC

Full list of author information is available at the end of the article

Background

Maize is known as corn which is widely grown in the Americas. In the United States alone, approximately 332 million metric tons of corn are grown annually [1]. Corn silk (*Stigmata Maydis*) is the female part of the corn. Although corn silk is often processed as agriculture waste, it is actually consumed as tea or regarded as an herb in traditional medicine. The main medicinal property of corn silk is to promote fluid excretion and reduce swelling [2]. In addition to its use as a natural diuretic, pharmacological studies of corn silk revealed antioxidant and anti-glycation activities that are used in diabetes, nephritis, or hypertension therapy [3]. In terms of chemical constituents related to these bio-activities, flavonoids such as luteolin, formononetin and apigenin are identified from *Stigmata Maydis* and their antioxidant properties have been illustrated [4–7]. Additionally, polysaccharides of *Stigmata Maydis* have been shown to reduce blood glucose and protein glycation in diabetic mice [8, 9].

Under physiological conditions, reactive oxygen species (ROS) and protein glycation are important molecules and essential biochemical events in the human body. However, excessive ROS or accumulated advanced glycation end-products (AGEs) may lead to tissue damage and aging [10]. For instance, diabetes is a complex metabolic disease with hyperglycaemia resulting from either insulin deficiency (Type 1 diabetes) or impaired insulin action and insulin secretory function (Type 2 diabetes) [11]. When diabetes is not well controlled, chronic hyperglycaemia leads to the progression of various diabetic complications via oxidative stress and AGE formation [12].

The loss of β -cell function and mass observed in uncontrolled diabetes may also be owing to glucotoxicity. Due to low levels of antioxidant enzymes, pancreatic β -cells are susceptible to oxidative stress via the production of excessive ROS under hyperglycaemic conditions [13]. As a result, glucotoxicity causes insulin secretory dysfunction and increased β -cell apoptosis, thus initiating a vicious cycle for glycaemic control [14, 15]. Moreover, low-level inflammation and additional oxidative stress from AGEs may affect β -cell proliferation and survival, thereby promoting β -cell failure [16].


Using corn silk constituents that possess both antioxidant and anti-glycation bio-activities, we aimed to generate a corn silk extract fraction that combines both activities to provide protective effects against glucotoxicity in insulin-secreting cells.

Methods

Extraction and partition of *Stigmata Maydis*

Stigmata Maydis was purchased from a Fu-Ji Chinese Traditional Medicine Store in Taipei on November 2008. Voucher specimens were deposited with the Herbarium

of the National Research Institute of Chinese Medicine (NHP-00351). The ethanol extract (YMS) was made by extracting the dried material of *Stigmata Maydis* (6 kg) with 100 L of 95 % ethanol at 55 °C for 7 h. Each fraction was made by partitioning the water suspended YMS (135 g) with corresponding solvents, such as *n*-hexane (Hex), ethyl acetate (EA), *n*-butanol (BuOH), to yield YMS-Hex, -BuOH, -EA, and -W fractions, respectively (Fig. 1a). Each YMS fraction and three flavonoid compounds, including apigenin (A), formononetin (F), and luteolin (L), were applied on a Merck thin-layer chromatography plate (Silica gel 60 F₂₅₄, 0.25 mm) (Darmstadt, Germany) under the development in a mobile phase of CHCl₃ and MeOH (8.5:1.5). Afterwards, separated spots were sprayed with anisaldehyde spray reagent and detected after ultraviolet absorption at 254 nm and 365 nm.



ABTS free radical scavenging assay

2,2'-azino-bis (3-ethylbenzthiazoline-6-sulfonic acid) (ABTS) was used for the measurement of antioxidant activity. Briefly, a reaction mix consisting of potassium persulfate (2.45 mM) in ABTS solution (7 mM) was prepared and kept in the dark at room temperature for at least 16 h before use. The intensively-coloured ABTS^{•+} solution was then diluted with 0.01 M phosphate buffered saline (PBS) to give a pH of 7.4 with an absorbance of 0.70 at 734 nm. The *Stigmata Maydis* fractions were diluted 100× with the ABTS^{•+} solution to a total volume of 1 ml. Absorbance was measured at 6 min after the addition of test reagents. A negative control was made with PBS instead of ABTS^{•+} solution. The % inhibitions by different concentrations of samples were calculated according to the following equation: $[1 - (\text{Abs}_{\text{sample} + \text{ABTS}^{\bullet+} \text{ solution}} / \text{Abs}_{\text{ABTS}^{\bullet+} \text{ solution}}) \times 100]$ [17].

Bovine serum albumin (BSA)-methylglyoxal (MG) assay and AGE preparation

This assay was used to evaluate protein glycation, and BSA fluorescence levels were measured. Briefly, BSA (10 mg/ml) was non-enzymatically glycated via incubation in 1 M PBS, pH 7.4, at 37 °C for 7 days in the presence of 1 mM MG and 3 mM sodium azide. The *Stigmata Maydis* fractions were tested at concentrations of 0.01, 0.02, 0.05, 0.1, and 1.0 mg/ml. Fluorescence of the samples was measured at the excitation and emission wavelengths of 335 and 385 nm, respectively, versus a blank containing the protein and MG. The % inhibition by different concentrations of samples was calculated according to the following equation: $[1 - (F_{\text{sample} + \text{BSA} + \text{glucose}} - F_{\text{sample} + \text{BSA}}) / (F_{\text{BSA} + \text{glucose}} - F_{\text{BSA}})] \times 100$. Aminoguanidine (AG) was used as a positive control.

The reactant under control condition was collected to generate AGEs through the dialysis and lyophilisation process. Products were kept at -80 °C for cell-based studies.

Cell culture

The clonal rat pancreatic β-cell line (BRIN-BD11) was kindly provided by prof. PR Flatt at University of Ulster, Coleraine, UK and routinely grown as a monolayer in culture dishes at 37 °C under 5 % CO₂/air with 90 % humidity. Cells were maintained in RPMI 1640 medium containing 10 % foetal bovine serum and 5 % penicillin and streptomycin mixture.

Cell viability assay (neutral red)

The cell viability assay was performed as previously described [18]. Briefly, at the end of cell treatments, the medium was replaced with the neutral red solution and incubated for another 2 h. Quantification of the uptake

of the neutral red by functional lysosomes in cells was spectrophotometrically measured at 540 nm.

Cell proliferation assay (WST-1)

The WST-1 cell proliferation assay was performed according to the manufacturer's protocol (Cayman Chemical). Briefly, cells were seeded on 96-well plates and the culture medium was replaced with various conditioned medium for 48 h. At the end of treatment, the WST-1 reagent was added and incubated for another 2 h. Finally, the plate was directly measured for absorbance at 450 nm.

Spectrofluorometric measurement of intracellular ROS

Intracellular ROS were measured by the CM-H₂DCFDA assay. Cells were cultured at 37 °C with various conditions which were described in figure legends. After 24 h, medium was replaced with the peroxide sensitive fluorescent probe, 5,6-dicarboxy-2,7-dichlorodihydro fluorescein diacetate (carboxy-H₂DCFDA; 20 μM), for an additional 30 min at 37 °C. The cells were then solubilised with 1 % SDS and 5 mM Tris HCl (pH 7.4). The fluorescence intensity of the lysate was determined using a spectrofluorometer with excitation and emission wavelengths of 495 nm and 517 nm, respectively.

Flow cytometry with annexin V/Propidium iodide (PI) staining

BRIN-BD11 cells were treated as mentioned above. Afterwards, they were trypsinised, pelleted, and resuspended in culture medium at a concentration of 1 × 10⁶ cells/ml. After transferring 0.5 ml of the cell suspension to a new tube, 10 μl media binding reagent and 1.25 μl annexin V-FITC were added. Following gentle vortexing, the mixture was incubated for 15 min at room temperature in the dark. After centrifuging at 1000xg for 5 min at room temperature, media was removed and 0.5 ml of cold 1× binding buffer and 10 μl propidium iodide were added. Following gentle vortexing, the sample was analysed on the flow cytometer within a 1 h period. The percentages of apoptotic and necrotic cells for each sample were estimated [19].

Gene expression analysis

BRIN-BD11 cells were seeded on 6 cm-dish (5 × 10⁵ cells/dish) and cultured under the condition described in Figure legends. At the end of experiments, total RNA were extracted and reverse transcribed. 50 ng of complementary (c)DNA of each sample was used for later polymerase chain reaction (PCR). Respective primer sequence, annealing temperature, and size of PCR product of each gene was listed below. Beta-

actin: For 5'-CGTAAAGACCTCTATGCCAA-3' and Rev 5'-AGCCATGCCAAATGTGTCAT-3'; 57 °C; 349b.p. Glucokinase: For 5'-AAGGGAACCTACATCGTAGGA-3' and Rev 5'-CATTGGCGGTCTTCATAGTA-3'; 57 °C; 130b.p. pancreatic and duodenal homeobox-1 (PDX-1): For 5'-CTCGCTGGGAACGCTGGAACA-3' and Rev 5'-GCTTTGGTGGATTTTCATCCACGG-3'; 55 °C; 225b.p. Insulin: For 5'-TGCCAGGCTTTTGTCAAACAGCACCTT-3' and Rev 5'-CTCCAGTGC CAAGGTCTGAA-3'; 52 °C; 187b.p. PCR products were separated by ethidium bromide stained gel electrophoresis, visualized, photographed with a digital camera, and quantified with Genetools 3.06 (Syngene, Frederick, MD, USA) [20].

Insulin secretion

BRIN-BD11 cells were plated on 24-well plates (0.5×10^5 cells/well) and incubated for 48 h with media containing 5.6 or 30.0 mM glucose. Then, after 1 h of pre-incubation with 1.1 mM glucose, cells were challenged either with 1.1 mM or with 16.7 mM glucose in Krebs-Ringer Bicarbonate Buffer for 20 min. The media were collected for insulin determination. Insulin concentrations were quantified by the Homogeneous Time-Resolved Fluorescence (HTRF) insulin assay and normalized to a million of total cell numbers [21].

Statistical analysis

Data were presented as mean \pm standard error of the mean. Statistical analyses were performed using GraphPad Prism (GraphPad, CA, USA). Single parameter-based comparisons were obtained from the unpaired student's *t*-test. *P* values less than 0.05 and 0.01 were considered to be significant. Multiparametric comparisons were performed using one-way ANOVA, followed by post-hoc analyses by Tukey's HSD protected least significant difference.

Results

Chemical characterization of four YMS fractions partitioned from an ethanolic crude extract of corn silk

As shown in Fig. 1b, all fractions and the three flavonoid compounds were individually developed on a single TLC plate. In addition, all samples were pooled and co-spotted on the same position on the 1st lane (Mix). According to the polarity of the solvents for extraction, the R_f values of all the detected compounds were distributed among the ranges of Hex (0.71 to 0.91), EA (0.52–0.71), BuOH (0.01 to 0.52) and H²O (0.0 to 0.01) fractions. The R_f values for two (apigenin and luteolin) of the three flavonoid compounds (apigenin: 0.66; formononetin: 0.74; luteolin: 0.57) were in the range of the YMS-EA fraction.

The ethyl acetate fraction (YMS-EA) most potently scavenges free radicals in vitro and protects against effects of H₂O₂ on β -cells

All fractions exhibited dose-dependent free radical scavenging effects (Fig. 1a). However, the effect of YMS-EA was superior to that of YMS-Hex, -BuOH, and -W (*p* < 0.001), as it provided 75 % inhibition at a concentration of 100 μ g/ml. Therefore, we chose YMS-EA as the major fraction and compared its effects with those of other test agents. First, to compare the protective effects of YMS fractions and reference drugs on H₂O₂-mediated ROS production and β -cell death, BRIN-BD11 cells were treated with H₂O₂ (125 μ M) in the presence of YMS fractions (100 μ g/ml), AG (2 mM), metformin (Met; 100 μ M), or trolox (Trox; 100 μ M) for 24 h.


As shown in Fig. 2b, there is nearly a three-fold increase in ROS levels in H₂O₂-treated BRIN-BD11 cells. The presence of YMS-EA significantly decreased ROS levels in H₂O₂-treated BRIN-BD11 cells. Comparing YMS-EA with other test agents, only YMS-Hex, YMS-W, and Trox exhibited similar activities. In addition, there was a 50 % reduction in the viability of H₂O₂-treated BRIN-BD11 cells after 24 h (Fig. 2c). YMS-EA significantly improved the cell viability of H₂O₂-treated BRIN-BD11 cells. YMS-Hex, YMS-W, AG, Met, and Trox provided similar protective effects.

YMS-EA and reference drugs attenuate the effects of acute H₂O₂ treatment on proliferation in BRIN-BD11 cells

In terms of the impact of acute H₂O₂ treatment on cell proliferation, BRIN-BD11 cells were transiently treated with H₂O₂ for 2 h, and proliferation was monitored at 24 h post-treatment. The cell proliferation rate of BRIN-BD11 cells treated with H₂O₂ (125 or 250 μ M) dropped over 50 % at 6 h post-treatment (Fig. 3a). At 24 h post-treatment, the cell proliferation ratio of BRIN-BD11 cells that were treated with the lower concentration of H₂O₂ (125 μ M) returned to original levels. After 48 h, there was a 50 % reduction in the proliferation of BRIN-BD11 cells that were treated with higher concentration of H₂O₂ (250 μ M). Therefore, the effects of YMS-EA and reference drugs were tested in the presence of 250 μ M H₂O₂.

After acute H₂O₂ challenge, the treatment with YMS-EA or reference drugs insufficiently improved cell proliferation (Fig. 3b). YMS-EA had a better effect at 50 μ g/ml compare with that at 100 μ g/ml. However, all of the reference drugs performed better than 100 μ g/ml YMS-EA (*p* < 0.001).

According to the cell death analysis, the transient treatment of BRIN-BD11 cells with H₂O₂ for 2 h for BRIN-BD11 cells caused late apoptosis in a dose-dependent manner at 24 h post-treatment. The apoptotic cell population reached 70 % after 24 h when



BRIN-BD11 cells were transiently challenged with 250 μM H₂O₂ (Fig. 3c). Under this condition, treatment with YMS-EA or reference drugs had no effect (Fig. 3d).

YMS-EA treatment ameliorates MG-mediated glycation, ROS production, and cell death

The anti-glycation activity was determined by the production efficiency of fluorescent AGEs, which were generated by co-incubating BSA and MG (Fig. 4a). All YMS fractions inhibited the formation of AGEs in a dose-dependent manner. At low concentrations (10 μg/ml), the inhibitory effect of YMS-EA was less effective than that of YMS-Hex and -BuOH. However, at high concentrations (1000 μg/ml), YMS-EA demonstrated the most effective anti-glycation activity when compared with YMS-Hex, -BuOH, and -W (*p* < 0.01, *p* < 0.001, and *p* < 0.001, respectively).


As shown in Fig. 4b, MG (800 μM)-treated BRIN-BD11 cells exhibited three-fold increase in ROS levels after 24 h. Treatment with 100 μg/ml YMS-EA significantly reduced ROS production. Similar effects were observed with YMS-W, Met, and Trox. AG appeared to be the most potent anti-glycation agent. Furthermore, after BRIN-BD11 cells were treated with 800 μM MG for 24 h, viability decreased by nearly 50 % (Fig. 4c). In the presence of YMS-EA, there was a significant increase in the viability of BRIN-BD11 cells. Similar effects were observed with YMS-W, Met, and Trox. It is important to note that AG treatment elicited a nearly 100 % protective effect in BRIN-BD11 cells.

YMS-EA treatment has no protective effects on AGE-mediated cell death and anti-proliferation in BRIN-BD11 cells

The presence of AGEs (3 mg/ml) in the culture medium significantly reduced viability at 48 h post-treatment (*p* < 0.05) (Fig. 5a). There was no difference in cell viability under this condition in the presence of YMS-EA (50 or 100 μg/ml). Additionally, treatment with 3 mg/ml AGEs had a potent anti-proliferation effect on BRIN-BD11 cells (*p* < 0.001). Surprisingly, instead of providing beneficial effects on cell proliferation, the presence of YMS-EA (100 μg/ml) significantly worsened cell proliferation (*p* < 0.05).

Addition of YMS-EA attenuates the hyperglycaemia-induced elevation of ROS, reduction of β-cell marker genes, and impairment of glucose responsiveness in BRIN-BD11 cells

Compared with cells cultured under 5.6 mM glucose, cells cultured under 30 mM glucose exhibited a 1.5-fold increase in ROS levels (Fig. 6a). The presence of YMS-EA significantly suppressed the induction of ROS by





30 mM glucose ($p < 0.01$). Both concentrations of YMS-EA were equally effective, whereas none of the reference drugs had ROS scavenging effects.

We further analysed the gene expression of β -cell markers in the presence of 5.6 mM or 30 mM glucose. There was a significant reduction in the mRNA levels of insulin, glucokinase, and pancreatic and duodenal homeobox-1 (PDX-1) (Fig. 6b). Consistently, the presence of YMS-EA attenuated this reduction in the mRNA levels of β -cell markers.


Finally, when BRIN-BD11 cells were cultured under 30 mM glucose for 48 h, the glucose responsiveness of

BRIN-BD11 cells was abolished (Fig. 6c). Treatment of 50 µg/ml YMS-EA restored insulin secretory activity in response to 16 mM glucose ($p < 0.01$). However, this beneficial effect did not appear when 100 µg/ml YMS-EA was used. Under the stimulated condition (16.7 mM glucose), the amount of insulin secreted from YMS-EA (50 µg/ml)-treated BRIN-BD11 cells was significantly more ($p < 0.01$) than that from YMS-EA (100 µg/ml)-treated cells. In contrast, in Fig. 6d, treatment of AG could only significantly enhanced insulin secretion under basal condition ($p < 0.05$) rather than stimulated condition.



Individual effects of three flavonoid compounds on dual activities and beta-cell protection

By employing previous experiments, individual effects of apigenin (A), formononetin (F) and luteolin (L) were examined. In Fig. 7a, only apigenin and luteolin provided a dose-dependent protective effect on H₂O₂ -induced β-cell death. In terms of anti-AGE formation, only apigenin at 100 µM can significantly inhibited AGE formation (*p* < 0.01). Furthermore, three flavonoids provided beta-cell protection against methylglyoxal. Formononetin appeared to be superior to other two (Fig. 7c). In terms of AGEs-inhibited cell proliferation, all flavonoids could provide beneficial effects. Instead, addition of apigenin and luteolin worsen the inhibitory effects of AGEs on cell proliferation. Finally, impaired glucose-responsiveness of hyperglycemia-damaged BRIN-BD11 cells was unable to be restored by three



flavonoids (Fig. 7d). However, addition of formononetin partially increased basal insulin secretion (Fig. 7e).

Discussion

Because a large amount of corn silk is treated as agriculture waste after the processing of corn, using a simple preparation process to develop a corn silk fraction with health benefits would promote the utilization of corn silk. In the present study, we aimed to make a fraction with dual bio-activities, including antioxidant and anti-glycation activities. Aminoguanidine, trolox, and metformin are three reference drugs with single or dual activities against oxidative stress and glycation [22–25]. Our in vitro results indicate that the YMS-EA fraction was the most effective fraction. Although the YMS-BuOH fraction was partitioned with *n*-butanol, which has the closest polarity index number compared to ethyl acetate, it exhibited significantly less bio-activity. The R_f

values of apigenin and luteolin were in the range of YMS-EA suggested that both flavonoids may contribute to the dual bioactivity of YMS-EA. Luteolin derivatives were also previously identified from EA fraction [3]. Interestingly, two independent research publications showing that apigenin has activity to scavenge ROS and luteolin intervenes the formation of AGEs support our view [6, 26].

The islets are known to have relatively low levels of antioxidants, and decreases in blood glutathione levels contribute to the accumulation of ROS in the islet during diabetes. Thus, the application of an antioxidation mechanism to protect beta-cell survival and function is a long-accepted concept [27]. The employment of antioxidants is an important strategy for the preservation of β -cell function, as shown in several experimental and clinical studies [28, 29]. Consistent with the above concept, our results demonstrate that YMS-EA possessed




Fig. 7 Evaluation of dual activities and beta-cell protective effects of three flavonoids. To explore the activity relationship between YMS-EA and three flavonoids, we repeated following experiments by using three flavonoids (μM). **a** H_2O_2 -induced cell death. Data are mean \pm SEM ($n = 6$). $***p < 0.001$ versus the None group. **b** AGE formation assay. Data are mean \pm SEM ($n = 3$). $***p < 0.001$ versus the None group. **c** Methylglyoxal-induced cell death. Data are mean \pm SEM ($n = 12$). $*p < 0.05$ and $***p < 0.001$ versus the None group. **d** AGEs inhibited cell proliferation assay. Data are mean \pm SEM ($n = 8$). $***p < 0.001$ versus the None group. **e** Glucose stimulated insulin secretion of hyperglycemia damaged cells. Data are mean \pm SEM ($n = 6$). $**p < 0.01$ versus the 1.1 mM glucose group-under the same culture condition. $^ap < 0.05$ and $^cp < 0.001$ versus the corresponding condition of None (30 mM Glucose) group

free radical scavenging activity and improved BRIN-BD11 cell viability by reducing ROS production in the presence of H_2O_2 . This activity of YMS-1 was better than that of YMS-BuOH, AG, and Met. In addition, YMS-EA attenuated the anti-proliferative effect of acute H_2O_2 treatment on BRIN-BD11 cells. However, the application of reference drugs appeared to be more effective than YMS-EA. In contrast, the presence of YMS-EA and other reference drugs could not effectively prevent BRIN-BD11 cells from previously triggered apoptosis by H_2O_2 . Therefore, the present study suggests that YMS-EA and reference drugs are more suitable for the prevention or intervention of ROS-induced cell death. Similar to reference drugs, YMS-EA could not rescue H_2O_2 -induced apoptotic cells. This was possibly because YMS-EA and reference drugs lack of the DNA repair activity against H_2O_2 -induced DNA damages and subsequent apoptosis [20, 30].

AGEs are generated from the nonenzymatic interaction between protein and carbohydrates and are

regarded as important mediators of diabetes-related complications. AGEs also play a role in beta-cell failure in diabetes [16]. MG is a reactive compound that is derived from glucose and fructose metabolism. It is not only is a ROS donor but plays a role in AGE formation [31]. In the current study, a high concentration of YMS-EA was the most effective at inhibiting MG-mediated AGE formation. However, only YMS-EA and YMS-W significantly inhibited MG-induced ROS production and improved cell survival in BRIN-BD11 cells. Among all test agents, AG was the most potent agent for the prevention of MG-mediated cell death. Our results also further indicate that YMS-EA could not provide any protection to enable β -cell survival against AGE-mediated toxicity. Our findings suggest that YMS-EA has a preventive, but not rescue, effect on the loss of β -cell mass and function in diabetes.

Finally, we evaluated whether YMS-EA could prevent glucotoxicity-induced β -cell dysfunction. The excessive entry of glucose has been shown to elicit β -cell injury

and increase the nonenzymatic glycation of cellular proteins in animal studies [32, 33]. A β -cell line cultured under high-glucose concentrations exhibits deteriorating outcomes in insulin, glucokinase, and PDX-1 expression [34–37]. Oxidative stress is also known to play an important role in high glucose-mediated β -cell dysfunction [38]. Interestingly, only YMS-EA could effectively reduce the level of ROS under high-glucose conditions. Such effect was associated with the restoration of the expression of important beta-cell marker genes. However, in terms of the insulin secretory function in response to glucose, the improvement in glucose responsiveness by 50 μ g/ml YMS-EA treatment disappeared when 100 μ g/ml YMS-EA was used. In the future, an optimized dosage for YMS-EA should be carefully examined.

By examining activities of three flavonoids, results actually pointed out similarity between the actions of three flavonoids and YMS-EA. Consistent with the indication that YMS-EA might contain apigenin and luteolin type of flavonoids, both compounds provided strong antioxidant effects while apigenin could also prevent AGE formation. Interestingly, apigenin, luteolin, and YMS-EA at high dose worsen AGEs-inhibited cell proliferation. As a result, the dual effects of YMS-EA might be derived from collaboration of those compounds.

Conclusions

In conclusion, our study provides some basis to support the notion that *Stigmata Maydis* could be developed as a dietary agent to protect β -cell survival and function against pathological oxidative stress and protein glycation in diabetes.

Abbreviations

ABTS: 2,2'-azino-bis (3-ethylbenzthiazoline-6-sulfonic acid); AGEs: Advanced glycation end-products; BSA: Bovine serum albumin; carboxy-H2DCFDA: 5,6-dicarboxy-2,7-dichlorodihydro fluorescein diacetate; HTRF: Homogeneous Time-Resolved Fluorescence; MG: Methylglyoxal; PBS: Phosphate buffered saline; PDX-1: Pancreatic and duodenal homeobox-1; PI: Propidium iodide; ROS: Reactive oxygen species

Acknowledgements

BRIN-BD11 cells were kindly provided by Prof. Flatt, P.R. from University of Ulster, UK. Manuscript was edited by a native English editor from biosciencewriters.

Funding

This work was funded by the National Research Institute of Chinese Medicine, Ministry of Health and Welfare (Grant No. MM10501-0274) and Cheng Hsin General Hospital (Grant Number: Chgh-103-42).

Availability of data and materials

The datasets supporting the conclusions of this article are included within the article.

Authors' contributions

CCC, WY, HYR, and JLC carried out the study. HKL and CCC designed the experiments and supervised the work. HJT helped to analyzed and interpreted the data. HKL, CCC, and WY wrote the manuscript. HCH and YCL helped to provide research data for revise manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Consent for publication

Not applicable in this section. This article is not a clinical study involving human participants.

Ethics approval and consent to participate

Not applicable in this section.

Author details

¹School of Pharmacy, College of Medicine, National Taiwan University, Taipei, Taiwan, ROC. ²Department of Psychiatry, Cheng Hsin General Hospital, Taipei, Taiwan, ROC. ³Department of Life Sciences and Institute of Genome Sciences, National Yang-Ming University, Taipei, Taiwan, ROC. ⁴Division of Basic Chinese Medicine, National Research Institute of Chinese Medicine, Ministry of Health and Welfare, Taipei, Taiwan, ROC. ⁵Department of Applied Science, National Hsinchu University of Education, Hsinchu, Taiwan, ROC. ⁶The Center of Translational Medicine, Taipei Medical University, Taipei, Taiwan, ROC. ⁷The Ph.D. Program for Medical Biotechnology, College of Medical Science and Technology, Taipei Medical University, Taipei, Taiwan, ROC. ⁸Institute of Neuroscience, Brain Research Centre, National Yang-Ming University, Taipei, Taiwan, ROC. ⁹Ph.D Program for the Clinical Drug Discovery from Herbal Medicine, College of Pharmacy, Taipei Medical University, Taipei, Taiwan, ROC.

Received: 27 May 2016 Accepted: 7 October 2016

Published online: 03 November 2016

References

- Nations FaAOotU. In: Division S, editor. Maize, rice and wheat: area harvested, production quantity, yield. 2009.
- Hasanudin K, Hashim P, Mustafa S. Corn silk (*Stigma maydis*) in healthcare: a phytochemical and pharmacological review. *Molecules* (Basel, Switzerland). 2012;17(8):9697–715.
- Ren SC, Ding XL. Study on determination methods of flavonoids from corn silk. *Food Sci*. 2004;25(3):139–42.
- Yu TT, Lu XX, Lian XJ, Zhang YQ. Composition analysis of flavonoids from corn silk with thin-layer chromatography and ultraviolet spectrophotometry. *Food Sci*. 2008;29(11):477–81.
- Silva MM, Santos MR, Caroco G, Rocha R, Justino G, Mira L. Structure-antioxidant activity relationships of flavonoids: a re-examination. *Free Radic Res*. 2002;36(11):1219–27.
- Wu CH, Yen GC. Inhibitory effect of naturally occurring flavonoids on the formation of advanced glycation endproducts. *J Agric Food Chem*. 2005;53(8):3167–73.
- Matsuda H, Wang T, Managi H, Yoshikawa M. Structural requirements of flavonoids for inhibition of protein glycation and radical scavenging activities. *Bioorg Med Chem*. 2003;11(24):5317–23.
- Zhao WZ, Yu ZP, Yu YD, Liu BQ, Liu JB. Research progress of polysaccharides from *Stigma maydis*. *Food Sci*. 2010;31(11):289–92.
- Suzuki R, Okada Y, Okuyama T. A new flavone C-glycoside from the style of *Zea mays* L. with glycation inhibitory activity. *Chem Pharm Bull* (Tokyo). 2003;51(10):1186–8.
- Stefano GB, Challenger S, Kream RM. Hyperglycemia-associated alterations in cellular signaling and dysregulated mitochondrial bioenergetics in human metabolic disorders. *Eur J Nutr*. 2016. doi:10.1007/s00394-016-1212-2.
- Lin Y, Sun Z. Current views on type 2 diabetes. *J Endocrinol*. 2010;204(1):1–11.
- Ajith TA, Vinodkumar P. Advanced glycation end products: association with the pathogenesis of diseases and the current therapeutic advances. *Curr Clin Pharmacol*. 2016;11(2):118–27.
- Brun T, Maechler P. Beta-cell mitochondrial carriers and the diabetogenic stress response. *Biochim Biophys Acta*. 2016;1863(10):2540–9.
- Kharroubi I, Ladriere L, Cardozo AK, Dogusan Z, Cnop M, Eizirik DL. Free fatty acids and cytokines induce pancreatic beta-cell apoptosis by different mechanisms: role of nuclear factor-kappaB and endoplasmic reticulum stress. *Endocrinology*. 2004;145(11):5087–96.
- Harmon JS, Stein R, Robertson RP. Oxidative stress-mediated, post-translational loss of MafA protein as a contributing mechanism to loss of

- insulin gene expression in glucotoxic beta cells. *J Biol Chem*. 2005;280(12):11107–13.
16. Lim M, Park L, Shin G, Hong H, Kang I, Park Y. Induction of apoptosis of beta cells of the pancreas by advanced glycation end-products, important mediators of chronic complications of diabetes mellitus. *Ann N Y Acad Sci*. 2008;1150:311–5.
 17. Boudouda HB, Zeghib A, Karioti A, Bilia AR, Ozturk M, Aouni M, Kabouche A, Kabouche Z. Antibacterial, antioxidant, anti-cholinesterase potential and flavonol glycosides of *Biscutella raphanifolia* (Brassicaceae). *Pak J Pharm Sci*. 2015;28(1):153–8.
 18. Chen CC, Hsu CY, Chen CY, Liu HK. Fructus Corni suppresses hepatic gluconeogenesis related gene transcription, enhances glucose responsiveness of pancreatic beta-cells, and prevents toxin induced beta-cell death. *J Ethnopharmacol*. 2008;117(3):483–90.
 19. Rieger AM, Nelson KL, Konowalchuk JD, Barreda DR. Modified annexin V/propidium iodide apoptosis assay for accurate assessment of cell death. *J Vis Exp*. 2011;(50):2597.
 20. Liu HK, McCluskey JT, McClellan NH, Flatt PR. Iterative exposure of clonal BRIN-BD11 cells to ninhydrin enables selection of robust toxin-resistant cells but with decreased gene expression of insulin secretory function. *Pancreas*. 2008;36(3):294–301.
 21. Stone VM, Dhayal S, Brocklehurst KJ, Lenaghan C, Sorhede Winzell M, Hammar M, Xu X, Smith DM, Morgan NG. GPR120 (FFAR4) is preferentially expressed in pancreatic delta cells and regulates somatostatin secretion from murine islets of Langerhans. *Diabetologia*. 2014;57(6):1182–91.
 22. Adeshara K, Tupe R. Antiglycation and cell protective actions of metformin and glipizide in erythrocytes and monocytes. *Mol Biol Rep*. 2016;43(3):195–205.
 23. Piro S, Rabuazzo AM, Renis M, Purrello F. Effects of metformin on oxidative stress, adenine nucleotides balance, and glucose-induced insulin release impaired by chronic free fatty acids exposure in rat pancreatic islets. *J Endocrinol Invest*. 2012;35(5):504–10.
 24. Thornalley PJ. Use of aminoguanidine (Pimagedine) to prevent the formation of advanced glycation endproducts. *Arch Biochem Biophys*. 2003;419(1):31–40.
 25. Sharma SS, Sayyed SG. Effects of trolox on nerve dysfunction, thermal hyperalgesia and oxidative stress in experimental diabetic neuropathy. *Clin Exp Pharmacol Physiol*. 2006;33(11):1022–8.
 26. Batra P, Sharma AK. Anti-cancer potential of flavonoids: recent trends and future perspectives. *3 Biotech*. 2013;3(6):439–59.
 27. Masuda Y, Vaziri ND, Li S, Le A, Hajjighasemi-Ossareh M, Robles L, Foster CE, Stamos MJ, Al-Abodullah I, Ricordi C, et al. The effect of Nrf2 pathway activation on human pancreatic islet cells. *PLoS One*. 2015;10(6):e0131012.
 28. Page KA, Reisman T. Interventions to preserve beta-cell function in the management and prevention of type 2 diabetes. *Curr Diab Rep*. 2013;13(2):252–60.
 29. Robertson RP, Harmon J, Tran PO, Tanaka Y, Takahashi H. Glucose toxicity in beta-cells: type 2 diabetes, good radicals gone bad, and the glutathione connection. *Diabetes*. 2003;52(3):581–7.
 30. Onaran I, Guven GS, Ozdas SB, Kanigur G, Vehid S. Metformin does not prevent DNA damage in lymphocytes despite its antioxidant properties against cumene hydroperoxide-induced oxidative stress. *Mutat Res*. 2006;611(1–2):1–8.
 31. Matafome P, Sena C, Seica R. Methylglyoxal, obesity, and diabetes. *Endocrine*. 2013;43(3):472–84.
 32. Tajiri Y, Moller C, Grill V. Long-term effects of aminoguanidine on insulin release and biosynthesis: evidence that the formation of advanced glycosylation end products inhibits B cell function. *Endocrinology*. 1997;138(1):273–80.
 33. Mullokandov EA, Franklin WA, Brownlee M. DNA damage by the glycation products of glyceraldehyde 3-phosphate and lysine. *Diabetologia*. 1994;37(2):145–9.
 34. Prentki M, Joly E, El-Assaad W, Roduit R. Malonyl-CoA signaling, lipid partitioning, and glucolipototoxicity: role in beta-cell adaptation and failure in the etiology of diabetes. *Diabetes*. 2002;51 Suppl 3:S405–13.
 35. Robertson RP, Zhang HJ, Pyzdrowski KL, Walseth TF. Preservation of insulin mRNA levels and insulin secretion in HIT cells by avoidance of chronic exposure to high glucose concentrations. *J Clin Invest*. 1992;90(2):320–5.
 36. Olson LK, Redmon JB, Towle HC, Robertson RP. Chronic exposure of HIT cells to high glucose concentrations paradoxically decreases insulin gene transcription and alters binding of insulin gene regulatory protein. *J Clin Invest*. 1993;92(1):514–9.
 37. Olson LK, Sharma A, Peshavaria M, Wright CV, Towle HC, Robertson RP, Stein R. Reduction of insulin gene transcription in HIT-T15 beta cells chronically exposed to a supraphysiologic glucose concentration is associated with loss of STF-1 transcription factor expression. *Proc Natl Acad Sci U S A*. 1995;92(20):9127–31.
 38. Tangvarasittichai S. Oxidative stress, insulin resistance, dyslipidemia and type 2 diabetes mellitus. *World J Diabetes*. 2015;6(3):456–80.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

