CORRECTION Open Access



Correction: Baicalin promotes apoptosis and inhibits proliferation and migration of hypoxia-induced pulmonary artery smooth muscle cells by up-regulating A2a receptor via the SDF-1/CXCR4 signaling pathway

Xiaoying Huang^{1*}, Wei Mao¹, Ting Zhang¹, Meibin Wang¹, Xuetao Wang¹, Yaozhe Li¹, Lin Zhang¹, Dan Yao¹, Xueding Cai¹ and Liangxing Wang^{1*}

Correction: BMC Complement Med Ther 18, 330 (2018) https://doi.org/10.1186/s12906-018-2364-9

Following publication of the original article [1], the authors identified an error in Fig. 7. The correct figure is given below.

The original article [1] has been corrected.

The online version of the original article can be found at https://doi.org/10.1186/s12906-018-2364-9.

*Correspondence:
Xiaoying Huang
zjwzhxy@126.com
Liangxing Wang
wzyxywlx@163.com

¹Division of Pulmonary Medicine, The First Affiliated Hospital of Wenzhou
Medical University, Key Laboratory of Heart and Lung,
Wenzhou, Zhejiang 325000, People's Republic of China



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.




Fig. 7 A2aR upregulation and Baicalin attenuated hypoxia induced PASMCs proliferation and migration. **a**: Cell density in each group as viewed under a microscope. **b**: CCK-8 values in each group. **c**: Migrated PASMCs in each group as viewed under a microscope. **d**: Cell migration rate in each group detected by Transwell assay. # P < 0.05, ##p < 0.01 vs. normoxia group, * P < 0.05 **P < 0.05 vs. hypoxia group; + P < 0.05, ++ P < 0.01. Comparison between groups. P < 0.05 = 3.

Published online: 18 April 2023

muscle cells by up-regulating A2a receptor via the SDF-1/CXCR4 signaling pathway. BMC Complement Altern Med. 2018;18:330. https://doi.org/10.1186/s12906-018-2364-9.

References

Huang X, Mao W, Zhang T, et al. Baicalin promotes apoptosis and inhibits proliferation and migration of hypoxia-induced pulmonary artery smooth

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.