

RESEARCH ARTICLE

Open Access

Identifying Chinese herbal medicine for premenstrual syndrome: implications from a nationwide database

Hsing-Yu Chen^{1,2,3†}, Ben-Shian Huang^{4,5†}, Yi-Hsuan Lin^{1,2,3}, Irene H Su⁶, Sien-Hung Yang^{1,3}, Jiun-Liang Chen^{1,3}, Jen-Wu Huang⁷ and Yu-Chun Chen^{5,8,9,10*}

Abstract

Background: Premenstrual syndrome (PMS) occurs in women during their reproductive age with a quite negative impact on their daily lives. Women with PMS experience a wide range of physical or psychological symptoms and seek treatment for them. Chinese herb medicine (CHM) is commonly used for PMS and the goal of this study is to investigate the prescription patterns of CHM for PMS by using a nationwide database.

Methods: Prescriptions of CHM were obtained from two million beneficiaries randomly sampled from the National Health Insurance Research Database, a nationwide database in Taiwan. The ICD-9 code 625.4 was used to identify patients with PMS. Association rule mining and social network analysis were used to explore both the combinations and the core treatments for PMS.

Results: During 1998-2011, a total of 14,312 CHM prescriptions for PMS were provided. Jia-Wei-Xiao-Yao-San (JWXYS) was the CHM which had the highest prevalence (37.5% of all prescriptions) and also the core of prescription network for PMS. For combination of two CHM, JWXYS with *Cyperus rotundus* L. was prescribed most frequently, 7.7% of all prescriptions, followed by JWXYS with *Leonurus heterophyllus* Sweet, 5.9%, and *Cyperus rotundus* L. with *Leonurus heterophyllus* Sweet, 5.6%.

Conclusions: JWXYS-centered CHM combinations were most commonly prescribed for PMS. To the best of our knowledge, this is the first pharmaco-epidemiological study to review CHM treatments for PMS. However, the efficacy and safety of these commonly used CHM were still lacking. The results of this study provide valuable references for further clinical trials and bench studies.

Keywords: Chinese herbal medicine, Premenstrual syndrome, Association rule mining, National health insurance research database, Social network analysis, Traditional Chinese medicine

Background

Premenstrual syndrome (PMS) is a common gynecological disease among women in reproductive age. Most women experienced one or more physical, psychological or behavioral symptoms preceding or during menses; for some women, these symptoms severely disrupt their normal life [1]. The prevalence of PMS is high due to the various symptoms of PMS, such as food cravings, abdominal

discomfort, headache, pain, depression, irritability and anxiety [2]. Around 75% of reproductive age women have mild symptoms and 20-30% of women have clinically significant PMS symptoms [3,4]. Additionally, 13-18% of women need treatment due to Premenstrual Dysphoric Disorder (PMDD), the severe form of PMS [5].

PMS symptoms have been attributed to fluctuations in levels of neurotransmitters in the central nervous system and/or cyclic hormonal changes. Central nervous system neurotransmitters such as serotonin (5-hydroxytryptamine) and γ -aminobutyric acid (GABA) appear to contribute to the pathogenesis of PMS [6-9]. Moreover, dynamic fluctuations in estradiol or progesterone in the late luteal phase

* Correspondence: yuchn.chen@googlemail.com

†Equal contributors

⁵School of Medicine, National Yang-Ming University, Taipei, Taiwan

⁸Department of Medical Research and Education, National Yang-Ming University Hospital, #152, Xin Min Rd, I-Lan 26042, Taiwan

Full list of author information is available at the end of the article

may be another possible mechanism underlying PMS symptoms [10,11]. Accordingly, selective serotonin reuptake inhibitors (SSRIs) are the mainstay treatment of PMS, and hormone therapy to suppress ovulation is often used as well. In addition, other psychoactive drugs, such as alprazolam or buspirone, have been used for PMS with less effect [8,12].

These pharmacologic approaches are associated with side effects that obviously concern women with PMS. As a result, compliance in women with PMS treated with SSRIs is rather poor due to adverse drug effects, such as headache, weight gain, drowsiness, sexual dysfunction, and insomnia [13,14]. As well, prescriptions of alprazolam are limited due to drug dependence, intolerance or even abuse [8]. For these reasons, various types of traditional Chinese medicine (TCM) treatments, including acupuncture and Chinese herbal medicine (CHM), are used by women as an integrative approach to PMS [15-17]. Although TCM use is increasing yearly, information about CHM prescriptions used for women with PMS is still lacking [18].

CHM prescriptions are part of a rather complicated system and every prescription is made according to each patient's TCM syndrome, or "Zheng" in Chinese, which is an overall summary of each patient's physical signs or symptoms. Different diseases have their specific preponderance of TCM syndrome and the TCM syndrome varies from patient to patient [19]. Based on individuals' diseases and TCM syndrome, different core treatments are prescribed and therefore it is feasible to discover the most important CHM and TCM understanding about a disease [20-22]. These findings are keys to further studies, especially new drug discovery, from the viewpoint of TCM system biology [23]. Application of association rule mining (ARM) and social network analysis (SNA) on a huge prescription database is a practical method for exploration of CHM core treatments and combinations, and several core treatments were successfully discovered in menopausal syndrome [24], primary dysmenorrhea [20], urticaria [21] and asthma [22].

The National Health Insurance Research Database (NHIRD), a nationwide health insurance claims database in Taiwan, is a comprehensive population-based tool that helpful to analyze the CHM prescription patterns [20-22,24]. The NHIRD is composed of all medical records, including ambulatory visits and hospitalization, reimbursed by the National Health Insurance (NHI) since 1995 [25]. Additionally, the NHI is unique in Taiwan, because both traditional Chinese medicine (CHM) and western medicine are equally reimbursed. Therefore, the option of using TCM or western medicine is completely unrestricted and largely dependent on the patient's preference or habits. The quality in reimbursement is crucial to eliminate the bias on intervention selection. By using the NHIRD, the

results of analysis on prescription patterns can be regarded as a consensus made by TCM doctors in Taiwan due to high coverage of the NHI, with over 99% of all Taiwanese population enrolled in the system [21,24].

We hypothesized that there should be core CHM treatment for PMS and this study aimed at discovering the core treatment patterns by applying ARM and SNA on the nationwide database. In addition to better understanding the viewpoint of TCM on PMS, further studies regarding efficacy, safety and mechanisms of action could be conducted based on the results of this work.

Methods

Data source

The NHIRD was used to obtain CHM prescriptions associated with the diagnosis of PMS. Two million individuals randomly sampled from the NHIRD and all medical claims of these individuals between 1998 and 2011 were extracted. This sample dataset is representative of the entire eligible Taiwanese population since no gender or age differences between sample and entire database were reported. The medical information contained in the NHIRD included patient's birth date, gender, visit date, hospitalization date, medical services providers, treatments, managements, and the reasons for visits or hospitalization. The International Classification of Disease 9th revision code (ICD-9) was used to identify the indications for medical services. Doctors may use up to three ICD-9 codes for every ambulatory visit and the first diagnosis was required to be the main reason for each visit. All TCM managements, including CHM, acupuncture and massage, were included.

Study subjects

To recognize the PMS patients, ICD-9 code 625.4 (Premenstrual tension syndromes) was used and subject selection was performed by Chen YC and Chen HY, independently. PMS was characterized by periodical affective and somatic disorders, and it was associated with various symptoms [26,27]. Other mood disorders or menstrual diseases should be excluded, such as dysmenorrhea, endometriosis, or perimenopausal diseases, to make a definitive diagnosis; therefore, ambulatory visits with any diagnosis other than PMS were excluded. Additionally, only visits with a single diagnosis of PMS were included to reduce the influence of co-morbidities since TCM doctors usually treat patients by modifying a patients' physical condition and by trying to treat all co-morbidities at once. Also, to eliminate the potential influence on choosing CHM, TCM visits with acupuncture and massage were excluded since these TCM modalities were thought to be adjuvant of CHM and may influence CHM prescription.

Study variables

Frequencies, durations and dosage of CHM contained in prescriptions were the primary variables in this study to identify single agents and combinations of CHM commonly used for PMS. Two types of CHM are reimbursed by the NHI, single herb (SH) and herbal formula (HF), which follow fixed proportions of SH according to TCM classics. Quality control of CHM is assured as SH and HF are all produced as concentrated powders by Good Manufacturing Practice (GMP) pharmaceutical factories. In clinical practice, TCM doctor usually use several HF and/or SH in one prescription according to patient's disease, TCM syndrome and associated symptoms. HF in prescriptions often serves as the core treatment for a disease and thus it is usually co-prescribed with other CHM according to TCM principles behind preparing prescriptions. The relationships between various CHM are typically quite complicated and a CHM network is commonly used to demonstrate the prescription patterns of CHM [20,28]. The frequency of connections among CHM is also assessed to verify the role of each CHM among network.

Statistical analysis

Descriptive statistics were used to explore the prescription patterns of CHM for PMS. The prescription frequency, presented as prevalence, of the most commonly used CHM and their combinations, was analyzed. Moreover, the efficacy and potential underlying mechanisms of the commonly used CHM were thoroughly searched and reviewed. The English, Latin, Chinese names or corresponding Kampo medicine names were used in combination with "premenstrual syndrome" or "premenstrual dysphoric disorder" as the key words in searching references. Pubmed and Medline were both used as the target databases and the last assessed date was March 13, 2014. Chen HY, Lin YH, Yang SH and Chen JL were responsible for reviewing the searched references and determined the mechanisms of each listed CHM.

To explore the combinations of CHM, ARM and SNA were applied to illustrate the relationships between commonly used CHM. ARM is a statistical model widely used in medical research to examine associations between drugs or disease conditions, such as CHM prescription patterns for menopausal syndrome and analysis on comorbidities of a disease [24,29]. Support and confidence factors are two key factors used in the ARM model to filter the significant associations among all CHM. Support factor was the prevalence of CHM combinations and confidence factor was similar to the conditional probability of a certain combination. For example, among two CHM, named CHM A and CHM B, support factor was presented as the probability ($CHM A \cap CHM B$) and confidence factor was presented as the probability ($CHM A \cap CHM B$)/

probability (CHM A); only if both support and confidence factor were higher than the pre-set threshold was the association "CHM A with CHM B" considered significant. Significant associations were interpreted as "if CHM A was prescribed, CHM B was commonly used as well as an adjuvant". Software R, version 2.15, and its statistical package "arule" was used to apply ARM model in this study. The threshold for support and confidence factors were set to 1% and 30%, and the threshold selected resulted from consensus among all co-authors of this work. The free-ware NodeXL of SNA (<http://nodexl.codeplex.com/>) was used to illustrate the associations between commonly used CHM for PMS.

Results

Demographic characteristics of TCM users

Among two million representative dataset, a total of 5,668 patients were identified as having visited TCM doctors for PMS. During 1998-2011, a total of 14,312 visits were made by TCM visitors, and CHM was the most commonly used modality of TCM, accounting for 96.6% of all visits (13,820 visits). Additionally, TCM doctors commonly prescribed multiple CHM concomitantly; 5.4 CHM, including SH and/or HF, were used in on average in one prescription. Additionally, CHM was usually prescribed three times a day (89.0% of all prescriptions), followed by four times a day (8.4%).

The most commonly used HF and SH

Among all prescriptions, Jia-Wei-Xiao-Yao-San (JWXYS) was the most commonly used CHM, accounting for 37.5% of all prescriptions, and it was also the most commonly used HF, followed by Dang-Gui-Shao-Yao-San (DGSYS) (2,064/14.9%) and Gui-Zhi-Fu-Ling-Wan (1,245/9.0%) (Table 1). Among single herbs, *Cyperus rotundus* L. was prescribed most frequently (2,485/18.0%), followed by *Leonurus heterophyllus* Sweet (2,097/15.2%) and *Corydalis yanhusuo* W. T. Wang (1,573/11.4%) (Table 2). Interestingly, although SH were usually used as an adjuvant to HF, *Cyperus rotundus* L. and *Leonurus heterophyllus* Sweet, both SH, were prescribed even more frequently than the HF DGSYS in this population. Moreover, the dose of commonly used HF was 2-3 times higher than SH, at approximately 4 gm/day versus less than 1.4 gm/day. (Tables 2 and 3).

Two CHM in combinations for PMS

JWXYS prescribed with *Cyperus rotundus* L. was the most frequently used two-CHM combination, accounting for 7.7% of all prescriptions (Table 3). JWXYS with *Leonurus heterophyllus* Sweet (818/5.9%) and *Cyperus rotundus* L. with *Leonurus heterophyllus* Sweet (773/5.6%) were the second and third the most frequently prescribed combinations. "HF with SH" were the most often used

Table 1 The top five most commonly used herbal formulas for premenstrual syndrome (N = 13,820)

Herbal formula	Constituents	Indication of TCM syndrome	Instances	Dose (gm/day)	Duration (day)	Prevalence (%)
Jia-Wei-Xiao-Yao-San (JWXYS)	<i>Paeonia lactiflora</i> Pall., <i>Bupleurum chinense</i> DC. DC., <i>Atractylodes macrocephala</i> Koidz., <i>Poria cocos</i> (Schw.) Wolf, <i>Angelica sinensis</i> (Oliv.) Diels., <i>Mentha haplocalyx</i> Briq., <i>Glycyrrhiza uralensis</i> Fisch., <i>Zingiber officinale</i> Rosc., <i>Paeonia suffruticosa</i> Andr., and <i>Gardenia jasminoides</i> Ellis.	Liver and spleen blood deficiency with heat transforming, liver qi stagnation	5,185	4.87	6.4	37.5
Dang-Gui-Shao-Yao-San	<i>Angelica sinensis</i> (Oliv.) Diels., <i>Ligusticum chuanxiong</i> Hort., <i>Paeonia lactiflora</i> Pall., <i>Atractylodes macrocephala</i> Koidz., <i>Poria cocos</i> (Schw.) Wolf, and <i>Alisma plantago-aquatica</i> L. var. <i>Alisma orientale</i> Samuels	Liver blood deficiency and disharmony of Liver and Spleen	2,064	4.34	6.4	14.9
Gui-Zhi-Fu-Ling-Wan	<i>Cinnamomum cassia</i> Blume, <i>Poria cocos</i> (Schw.) Wolf, <i>Paeonia lactiflora</i> Pall., <i>Paeonia suffruticosa</i> Andr., and <i>Prunus persica</i> (L.) Batsch (or <i>Prunus davidiana</i> (Carr.) Franch.)	Blood stasis	1,245	3.87	6.3	9.0
Wen-Jing-Tang	<i>Evodia rutaecarpa</i> (Juss.) Benth., (or <i>Evodia rutaecarpa</i> (Juss.) Benth. var. <i>officinalis</i> (Dode) Huang, <i>Evodia rutaecarpa</i> (Juss.) Benth. var. <i>bodinieri</i> (Dode) Huang), <i>Cinnamomum cassia</i> Blume, <i>Angelica sinensis</i> (Oliv.) Diels., <i>Paeonia lactiflora</i> Pall., <i>Ligusticum chuanxiong</i> Hort., <i>Panax ginseng</i> C. A. Meyer, <i>Glycyrrhiza uralensis</i> Fisch., <i>Equus asinus</i> L., <i>Ophiopogon japonicus</i> Ker-Gawl., <i>Pinellia ternata</i> (Thunb.) Breit, and <i>Zingiber officinale</i> Rosc.	Blood stasis and deficient cold	1,120	4.25	6.3	8.1
Shao-Fu-Zhu-Yu-Tang	<i>Foeniculum vulgare</i> Mill., <i>Zingiber officinale</i> Rosc., <i>Cinnamomum cassia</i> Blume, <i>Angelica sinensis</i> (Oliv.) Diels., <i>Ligusticum chuanxiong</i> Hort., <i>Paeonia lactiflora</i> Pall. or <i>Paeonia veitchii</i> Lynch, <i>Trogopterus xanthipes</i> Milne-Edwards or <i>Pteromys volans</i> L.	Qi stagnation and blood stasis	994	3.97	6.2	7.2

combinations (80%), rather than “SH with SH” (20%) concept in TCM.

Core treatment for PMS

JWXYS was the core treatment for PMS because it accounted for the highest prevalence among all CHM and 80% of the most commonly used two-CHM combinations. Additionally, the central role of JWXYS can be found in the network of commonly used CHM (Figure 1). Other CHM scattered peripherally represented their adjuvant action of PMS.

Discussion

The key finding of this study is that JWXYS was the core treatment for PMS and certain CHM, such as *Cyperus rotundus* L. and *Leonurus heterophyllus* Sweet, were frequently prescribed with JWXYS. This is the first pharmaco-epidemiological study about CHM for PMS. Among all CHM used for PMS, only JWXYS was examined for its efficacy and included in systematic review [15,17]. The effectiveness and safety of other commonly used CHM, especially when they are combined with JWXYS, have not been studied yet, and CHM included in

Table 2 The top ten most commonly used single herb for premenstrual syndrome (N = 13,820)

Latin name	Indication of TCM syndrome	Instances	Dose (gm/day)	Duration (day)	Prevalence (%)
<i>Cyperus rotundus</i> L.	Qi stagnation in liver	2,485	1.38	6.3	18.0
<i>Leonurus heterophyllus</i> Sweet	Blood stasis	2,097	1.33	6.3	15.2
<i>Corydalis yanhusuo</i> W. T. Wang	Qi stagnation and blood stasis	1,573	1.28	6.2	11.4
<i>Salvia miltiorrhiza</i> Bge.	Blood stasis	1,281	1.30	6.4	9.3
<i>Eucommia ulmoides</i> Oliv.	Kidney and liver deficiency	1,023	1.26	6.3	7.4
<i>Scutellaria baicalensis</i> Georgi	Dampness-heat	858	1.30	6.5	6.2
<i>Dipsacus asperoides</i> C. Y. Cheng at T. M. Ai	Blood stasis	644	1.17	6.4	4.7
<i>Cuscuta chinensis</i> Lam., or <i>Cuscuta japonica</i> Choisy	Kidney yang and yin deficiency	639	1.44	6.6	4.6
<i>Pueraria lobata</i> (Willd.) Ohwi or <i>Pueraria thomsonii</i> Benth.	External wind, fluid deficiency	616	1.33	6.4	4.5
<i>Paeonia suffruticosa</i> Andr.	Blood stasis and heat	605	1.43	6.6	4.4

Table 3 The top ten most commonly used two-herb combinations of Chinese herbal medicine (CHM) for premenstrual syndrome (N = 13,820)

CHM A		CHM B	Instances	Prevalence (%)
JWXYS	with	<i>Cyperus rotundus</i> L.	1,059	7.7
JWXYS	with	<i>Leonurus heterophyllus</i> Sweet	818	5.9
<i>Cyperus rotundus</i> L.	with	<i>Leonurus heterophyllus</i> Sweet	773	5.6
JWXYS	with	<i>Corydalis yanhusuo</i> W. T. Wang	564	4.1
<i>Cyperus rotundus</i> L.	with	<i>Corydalis yanhusuo</i> W. T. Wang	512	3.7
JWXYS	with	<i>Salvia miltiorrhiza</i> Bge.	467	3.4
<i>Cyperus rotundus</i> L.	with	<i>Salvia miltiorrhiza</i> Bge.	377	2.7
JWXYS	with	Gui-Zhi-Fu-Ling-Wan	375	2.7
JWXYS	with	<i>Scutellaria baicalensis</i> Georgi	262	1.9
JWXYS	with	<i>Curcuma longa</i> L. or <i>Curcuma aromatica</i> Salisb.	251	1.8

latest systematic review are not exactly commonly used in daily practice [17]. The results of this study provide ideal candidate for further studies as we strive to achieve the goal of establishing evidence-based medicine viewpoints on CHM.

JWXYS has been reported to have anxiolytic and anti-depression effects in animal models although its effectiveness on PMS remains unclear (Table 4). From TCM's

viewpoint, JWXYS, which was developed about a thousand years ago in the TCM classic of prescriptions, is used to treat liver and spleen blood deficiency with heat transforming, which fairly represents mood disorder, gastrointestinal dysfunction and menstrual problems due to endocrine disorder. Therefore, it is frequently used to treat primary dysmenorrhea [20], menopausal syndrome [24], depression [30] and functional dyspepsia [31]. The high




Table 4 Possible pharmacological mechanisms of Chinese herbal medicine (CHM) used for premenstrual syndrome

Name	Ingredients	Possible mechanisms of extract of SH
Single herb (SH)		
<i>Cyperus rotundus</i> L.	Hydroalcoholic extract	Anti-oxidative effect [33]
	Essential oil	Anti-oxidative effect [34]
<i>Leonurus heterophyllus</i> Sweet	Methanol extract	Anti-oxidative effect [35]
<i>Corydalis yanhusuo</i> W. T. Wang	Tetrahydropalmatine	Anxiolytic effect [36]
		Analgesic effect [37]
<i>Salvia miltiorrhiza</i> Bge.	Miltirone	Anxiolytic effect [38]
	Salvianolic acid B	Anti-depressant effect [39]
		Analgesic effect [40]
		Anti-oxidative effect [41]
	Tanshinone IIA	Analgesic effect [42]
<i>Eucommia ulmoides</i> Oliv.	Water extract	Anti-oxidative effect [43,44]
Herbal formula (HF)		
Jia-Wei-Xiao-Yao-San (JWSYS)		Anti-depressant effect on animal model [45,46]
		IL-6 related anti-depressant effect on menopausal women with mood disorder [47]
		Anxiolytic effect on animal model [48]
		Reduced proinflammatory cytokine [49]
Dang-Gui-Shao-Yao-San		Anti-depressant effect on animal model [50]
		Anti-oxidative effect in vitro [51]
Wen-Jing-Tang		Anti-depressant effect for menopausal women [52]

prevalence of psychological discomfort and gastrointestinal problems among PMS women in Taiwan may be the reason for the leading role of JWXYS in all CHM [32].

In addition to this core treatment, several CHM were found to be significantly associated with core treatments, and thus a network of CHM prescribed for PMS can be established (Figure 1). These combinations of CHM were valuable guides for clinical practice since TCM doctors usually use several CHM in one prescription (5.4 CHM on average). This finding was expected as it has its foundation in TCM teachings and with potential pharmacological effects of CHM, CHM prescriptions are made according to an individual's constitution, namely *bian-zheng-lun-zhi*, which means the identification of TCM syndrome and treatment. It is reasonable since human physiology is complex and multiple strategies to treat PMS may be needed. Interestingly, anti-oxidative effects are reported extensively among the CHM surrounding JWXYS [33,35,41,43], which may complement JWXYS since JWXYS lacks anti-oxidative effect. The anti-oxidative effects exclusively found in SH strongly imply the importance of oxidative stress on PMS, such as *Cyperus rotundus* L. and *Leonurus heterophyllus* Sweet (Table 4). To date, only one study based on 41 patients discussed the potential association between PMS and this imbalance of oxidants/anti-oxidants [53]. Thus, the essential role of anti-oxidation therapy for PMS will require further

studies in order to be defined properly. This finding represents the merit of using ARM and social network analysis to analyze the relations between CHM, and not just focusing on the use of a single CHM.

Furthermore, same treatments for different diseases, another feature of CHM prescription, could be found when comparing the prescription patterns among PMS, primary dysmenorrhea and menopausal syndrome [20]. For example, JWSYS was also commonly used for mood disorder in menopausal women [24] and DGSYS was frequently used for primary dysmenorrhea and PMS [20]. This concept is quite different from western medicine which often treats PMS after excluding other diagnoses such as dysmenorrhea. Therefore, the treatment principle considerably differs: analgesia for primary dysmenorrhea and mood modifying for PMS. On TCM's viewpoint, prescriptions are made not only according to "disease" but also TCM syndrome, "zheng", and therefore it is not uncommon to prescribe same formula to various diseases if the same TCM syndrome is diagnosed [54].

"Qi stagnation and blood stasis" seemed the most prevalent TCM syndrome for PMS. Although it was not feasible to perform detailed analysis on TCM syndrome of PMS since TCM syndrome was not recorded in the NHIRD, the distribution of TCM syndrome could be shown when summarizing the indications of commonly used CHM [21,55]. Ten out of top 15 commonly used

CHM are indicated to manage “qi stagnation” or “blood stasis” (Tables 1 and 2) and JWXYS is effective in affective disorder caused by liver qi stagnation syndrome [56]. Qi stagnation usually represented lesion distension or pain due to impeded qi circulation and mood disorders, such as depression or anxiety and depression was highly associated with liver qi stagnation [57]. Further, severe qi stagnation may obstruct blood flow to cause blood stasis, but blood stasis may aggravate qi stagnation conversely. This TCM syndrome and pathogenesis reflected the concept about PMS: mood disorders (qi stagnation) and menstrual discomfort (blood stasis) are highly related. For this reason, regulation on menstrual cycle is needed to treat PMS for TCM doctors and that is why many CHM other than JWXYS are added into one prescription for PMS.

This study explored the core formula, JWXYS, and important combinations of CHM for PMS. This result is especially crucial in searching for candidates that are suitable for clinical or bench studies for PMS since CHM prescriptions are usually different from textbooks themselves [58]. The merit of this study is that the subject selection is done from a nationwide database and thus possible selection and referral bias can be eliminated. Nonetheless, there are a number of limitations to consider. First, the counts of prescriptions for PMS might be underestimated since only visits with a single diagnosis of PMS were used to identify the study population in this study. Nevertheless, it is believed that the use of a single diagnosis is helpful in eliminating unnecessary bias, especially when PMS is a syndrome with various presentations and the impact from co-morbidities may be considerable. Second, the validity of ICD-9 code for the diagnosis of PMS should be considered. In Taiwan, the first diagnosis code of each ambulatory visit represents the main reason for the visit. As physicians and institutions are penalized for fraudulent coding, monthly internal monitoring are usually performed to validate diagnoses. This internal monitoring theoretically improves the accuracy of the database. Therefore, we believe the diagnostic codes are specific and the registration bias can be minimized. Third, the efficacy of therapies was not determined in this study. Despite the fact that the nationwide database contains large amounts of prescription data, there are no data on PMS severities, laboratory or image examinations since these data are not required in the electronic medical records. Consequently, the effectiveness of CHM for PMS is unable to be assessed by using this database.

Conclusions

The high prevalence and role of the core treatment JWXYS for PMS is shown in this study. Also, the combinations of CHM with JWXYS, such as *Cyperus rotundus* L. and *Leonurus heterophyllus* Sweet, demonstrate the potential effectiveness of anti-oxidant agents on PMS.

Based on these results, further studies about CHM for PMS are warranted both in bench research and clinical trials to examine the effectiveness of these therapies.

Ethical approval

This study was approved by the Institutional Review Board (IRB) of Chang Gung Memorial Foundation, Taipei, Taiwan (IRB: 101-3604B).

Abbreviations

PMS: Premenstrual syndrome; CHM: Chinese herbal medicine, (JWXYS) Jia-Wei-Xiao-Yao-San; ICD-9: The International classification of disease, 9th revision; PMDD: Premenstrual dysphoric disorder; GABA: γ -aminobutyric acid; ARM: Association rule mining; NHIRD: The National Health Insurance Research Database; NHI: The National Health Insurance; SNA: Social network analysis; TCM: Traditional Chinese medicine.

Competing interests

The authors report that they have no conflicts of interest to declare.

Authors' contributions

HC and BH were responsible for study design, result interpretation and drafted the manuscript. HC also participated in data management. IS revised the manuscript and provided expert opinions as a professional gynecologist. YL, SY, JC, and JH participated in study design and result interpretation. In addition, HC, YL, SY and JC were the investigators to review and summarize the pharmacologic mechanisms of CHM. YC participated in data management, statistics analysis and helped to manuscript writing. All authors read and approved the final version of this manuscript.

Authors' information

HC, YL, SY and JC are TCM doctors of Division of Chinese Internal Medicine, Center for Traditional Chinese Medicine, Chang Gung Memorial Hospital. In Taiwan, it has one of the largest TCM departments of all medical centers. HC is responsible for teaching evidence-based medicine of TCM in this department. By collaborating with YC, who is the head of Department of Medical Research and Education, National Yang-Ming University Hospital, several epidemiology studies about TCM are conducted. We are interested in how and why CHM are prescribed for patients for individual diseases and we believe this information is critical for TCM on the viewpoints of evidence-based medicine. These results are beneficial to study candidate selection and results of further clinical trials or bench studies will be much more adherent to clinical practice.

Acknowledgements

We thank the data management department of the Bureau of National Health Insurance for data retrieval. All the interpretations and conclusions in this study do not represent the opinion of this institute.

Author details

¹Division of Chinese Internal Medicine, Center for Traditional Chinese Medicine, Chang Gung Memorial Hospital, Taoyuan, Taiwan. ²Graduate Institute of Clinical Medical Sciences, College of Medicine, Chang Gung University, Taoyuan, Taiwan. ³School of Traditional Chinese Medicine, College of Medicine, Chang Gung University, Taoyuan, Taiwan. ⁴Department of Obstetrics and Gynecology, National Yang-Ming University Hospital, I-Lan, Taiwan. ⁵School of Medicine, National Yang-Ming University, Taipei, Taiwan. ⁶Department of Reproductive Medicine, University of California, La Jolla, San Diego, CA, USA. ⁷Department of Surgery, National Yang-Ming University Hospital, I-Lan, Taiwan. ⁸Department of Medical Research and Education, National Yang-Ming University Hospital, #152, Xin Min Rd, I-Lan 26042, Taiwan. ⁹Faculty of Medicine, School of Medicine, National Yang-Ming University, Taipei, Taiwan. ¹⁰Institute of Hospital and Health Care Administration, School of Medicine, National Yang-Ming University, Taipei, Taiwan.

Received: 27 March 2014 Accepted: 24 June 2014

Published: 27 June 2014

References

- Yonkers KA, O'Brien PM, Eriksson E: **Premenstrual syndrome.** *Lancet* 2008, **371**:1200–1210.
- Hartlage SA, Freels S, Gotman N, Yonkers K: **Criteria for premenstrual dysphoric disorder: secondary analyses of relevant data sets.** *Arch Gen Psychiatry* 2012, **69**:300–305.
- Campbell EM, Peterkin D, O'Grady K, Sanson-Fisher R: **Premenstrual symptoms in general practice patients. Prevalence and treatment.** *J Reprod Med* 1997, **42**:637–646.
- Borenstein J, Chiou CF, Dean B, Wong J, Wade S: **Estimating direct and indirect costs of premenstrual syndrome.** *J Occup Environ Med* 2005, **47**:26–33.
- Hälbreich U, Borenstein J, Pearlstein T, Kahn LS: **The prevalence, impairment, impact, and burden of premenstrual dysphoric disorder (PMS/PMDD).** *Psychoneuroendocrinology* 2003, **28**(Suppl 3):1–23.
- Rapkin AJ, Winer SA: **Premenstrual syndrome and premenstrual dysphoric disorder: quality of life and burden of illness.** *Expert Rev Pharmacoecon Outcomes Res* 2009, **9**:157–170.
- Rapkin AJ, Edelmann E, Chang LC, Reading AE, McGuire MT, Su TP: **Whole-blood serotonin in premenstrual syndrome.** *Obstet Gynecol* 1987, **70**:533–537.
- Freeman EW, Rickels K, Sondheimer SJ, Polansky M: **A double-blind trial of oral progesterone, alprazolam, and placebo in treatment of severe premenstrual syndrome.** *JAMA* 1995, **274**:51–57.
- Smith MJ, Adams LF, Schmidt PJ, Rubinow DR, Wassermann EM: **Abnormal luteal phase excitability of the motor cortex in women with premenstrual syndrome.** *Biol Psychiatry* 2003, **54**:757–762.
- Schmidt PJ, Nieman LK, Danaceau MA, Adams LF, Rubinow DR: **Differential behavioral effects of gonadal steroids in women with and in those without premenstrual syndrome.** *N Engl J Med* 1998, **338**:209–216.
- Wyatt K, Dimmock P, Jones P, Obhrai M, O'Brien S: **Efficacy of progesterone and progestogens in management of premenstrual syndrome: systematic review.** *BMJ* 2001, **323**:776–780.
- Rickels K, Freeman E, Sondheimer S: **Buspirone in treatment of premenstrual syndrome.** *Lancet* 1989, **1**:777.
- Marjoribanks J, Brown J, O'Brien PMS, Wyatt K: **Selective serotonin reuptake inhibitors for premenstrual syndrome.** *Cochrane Database Syst Rev* 2013(6):CD001396. doi:10.1002/14651858.CD001396.pub3.
- Hu XH, Bull SA, Hunkeler EM, Ming E, Lee JY, Fireman B, Markson LE: **Incidence and duration of side effects and those rated as bothersome with selective serotonin reuptake inhibitor treatment for depression: patient report versus physician estimate.** *J Clin Psychiatry* 2004, **65**:959–965.
- Jing Z, Yang X, Ismail KMK, Chen XY, Wu T: **Chinese herbal medicine for premenstrual syndrome.** *Cochrane Database Syst Rev* 2009(1):CD006414. doi:10.1002/14651858.CD006414.pub2.
- Kim SY, Park HJ, Lee H, Lee H: **Acupuncture for premenstrual syndrome: a systematic review and meta-analysis of randomised controlled trials.** *BJOG* 2011, **118**:899–915.
- Jang S, Kim D, Choi MS: **Effects and treatment methods of acupuncture and herbal medicine for premenstrual syndrome/premenstrual dysphoric disorder: systematic review.** *BMC Complement Altern Med* 2014, **14**:11.
- Chang LC, Huang N, Chou YJ, Lee CH, Kao FY, Huang YT: **Utilization patterns of Chinese medicine and Western medicine under the National Health Insurance Program in Taiwan, a population-based study from 1997 to 2003.** *BMC Health Serv Res* 2008, **8**:170.
- Yang SH, Chen HY, Lin YH, Chen YC: **The Exploration of Disease Pattern, Zheng, for Differentiation of Allergic Rhinitis in Traditional Chinese Medicine Practice.** *Evid Based Complement Alternat Med* 2012, **2012**:7.
- Chen HY, Lin YH, Su IH, Chen YC, Yang SH, Chen JL: **Investigation on Chinese herbal medicine for primary dysmenorrhea: implication from a nationwide prescription database in Taiwan.** *Complement Ther Med* 2014, **22**:116–125.
- Lin YH, Chen YC, Hu S, Chen HY, Chen JL, Yang SH: **Identifying core herbal treatments for urticaria using Taiwan's nationwide prescription database.** *J Ethnopharmacol* 2013, **148**:556–562.
- Chen HY, Lin YH, Thien PF, Chang SC, Chen YC, Lo SS, Yang SH, Chen JL: **Identifying core herbal treatments for children with asthma: implication from a Chinese herbal medicine database in Taiwan.** *Evid Based Complement Alternat Med* 2013, **2013**:125943.
- Li J, Lu C, Jiang M, Niu X, Guo H, Li L, Bian Z, Lin N, Lu A: **Traditional Chinese medicine-based network pharmacology could lead to new multicomponent drug discovery.** *Evid Based Complement Alternat Med* 2012, **2012**:149762.
- Chen HY, Lin YH, Wu JC, Chen YC, Yang SH, Chen JL, Chen TJ: **Prescription patterns of Chinese herbal products for menopausal syndrome: analysis of a nationwide prescription database.** *J Ethnopharmacol* 2011, **137**:1261–1266.
- Chen HY, Lin YH, Wu JC, Chen YC, Thien PF, Chen TJ, Yang SH, Chen JL, Lo SS: **Characteristics of pediatric traditional Chinese medicine users in Taiwan: a nationwide cohort study.** *Pediatrics* 2012, **129**:e1485–1492.
- Dickerson LM, Mazyck PJ, Hunter MH: **Premenstrual syndrome.** *Am Fam Physician* 2003, **67**:1743–1752.
- Johnson SR: **Premenstrual syndrome, premenstrual dysphoric disorder, and beyond: a clinical primer for practitioners.** *Obstet Gynecol* 2004, **104**:845–859.
- Gao Y, Liu Z, Wang CJ, Fan XS, Xie JY: **Chinese medicine formula network analysis for core herbal discovery.** *Brain Informatics* 2012, **7670**:255–264.
- Tai YM, Chiu HW: **Comorbidity study of ADHD: applying association rule mining (ARM) to National Health Insurance Database of Taiwan.** *Int J Med Inf* 2009, **78**:e75–83.
- Zhang Y, Han M, Liu Z, Wang J, He Q, Liu J: **Chinese herbal formula xiao yao san for treatment of depression: a systematic review of randomized controlled trials.** *Evid Based Complement Alternat Med* 2012, **2012**:931636.
- Qin F, Huang X, Ren P: **Chinese herbal medicine modified xiaoyao san for functional dyspepsia: meta-analysis of randomized controlled trials.** *J Gastroenterol Hepatol* 2009, **24**:1320–1325.
- Hsiao MC, Liu CY, Chen KC, Hsieh TT: **Characteristics of women seeking treatment for premenstrual syndrome in Taiwan.** *Acta Psychiatr Scand* 2002, **106**:150–155.
- Yazdanparast R, Ardestani A: **In vitro antioxidant and free radical scavenging activity of *Cyperus rotundus*.** *J Med Food* 2007, **10**:667–674.
- Kilani S, Ledauphin J, Bouhler I, Ben Sghaier M, Boubaker J, Skandrani I, Mosrati R, Ghedira K, Barillier D, Chkir-Ghedira L: **Comparative study of *Cyperus rotundus* essential oil by a modified GC/MS analysis method. Evaluation of its antioxidant, cytotoxic, and apoptotic effects.** *Chem Biodivers* 2008, **5**:729–742.
- Matkowsky A, Piotrowska M: **Antioxidant and free radical scavenging activities of some medicinal plants from the Lamiaceae.** *Fitoterapia* 2006, **77**:346–353.
- Henkes H, Franz M, Kendall O, Monroe J, Legaspi A, LeDoux J, Haese C, Williams D, McCall S, Johnson AD, Ceremuga TE: **Evaluation of the anxiolytic properties of tetrahydropalmatine, a *Corydalis yanhusuo* compound, in the male Sprague-Dawley rat.** *AANA J* 2011, **79**:S75–80.
- Cao FL, Shang GW, Wang Y, Yang F, Li CL, Chen J: **Antinociceptive effects of intragastric dl-tetrahydropalmatine on visceral and somatic persistent nociception and pain hypersensitivity in rats.** *Pharmacol Biochem Behav* 2011, **100**:199–204.
- Imanshahidi M, Hosseinzadeh H: **The pharmacological effects of *Salvia* species on the central nervous system.** *Phytother Res* 2006, **20**:427–437.
- Feng Y, You Z, Yan S, He G, Chen Y, Gou X, Peng C: **Antidepressant-like effects of salvianolic acid B in the mouse forced swim and tail suspension tests.** *Life Sci* 2012, **90**:1010–1014.
- Isacchi B, Fabbri V, Galeotti N, Bergonzi MC, Karioti A, Ghelardini C, Vannucchi MG, Bilia AR: **Salvianolic acid B and its liposomal formulations: anti-hyperalgesic activity in the treatment of neuropathic pain.** *Eur J Pharm Sci* 2011, **44**:552–558.
- Zhang YA, Li X, Wang ZZ: **Antioxidant activities of leaf extract of *Salvia miltiorrhiza* Bunge and related phenolic constituents.** *Food Chem Toxicol* 2010, **48**:2656–2662.
- Sun S, Yin Y, Yin X, Cao F, Luo D, Zhang T, Li Y, Ni L: **Anti-nociceptive effects of Tanshinone IIA (TIIA) in a rat model of complete Freund's adjuvant (CFA)-induced inflammatory pain.** *Brain Res Bull* 2012, **88**:581–588.
- Luo J, Tian C, Xu J, Sun Y: **Studies on the antioxidant activity and phenolic compounds of enzyme-assisted water extracts from *Du-zhong* (*Eucommia ulmoides* Oliv.) leaves.** *J Enzyme Inhib Med Chem* 2009, **24**:1280–1287.
- Park SA, Choi MS, Jung UJ, Kim MJ, Kim DJ, Park HM, Park YB, Lee MK: ***Eucommia ulmoides* Oliver leaf extract increases endogenous antioxidant activity in type 2 diabetic mice.** *J Med Food* 2006, **9**:474–479.

45. Park SW, Kim YK, Lee JG, Kim SH, Kim JM, Yoon JS, Park YK, Lee YK, Kim YH: **Antidepressant-like effects of the traditional Chinese medicine kami-shoyo-san in rats.** *Psychiatry Clin Neurosci* 2007, **61**:401–406.
46. Oh JK, Kim Y-S, Park H-J, Lim E-M, Pyun K-H, Shim I: **Antidepressant effects of Soyo-san on immobilization stress in ovariectomized female rats.** *Biol Pharm Bull* 2007, **30**:1422–1426.
47. Yasui T, Yamada M, Uemura H, Ueno S, Numata S, Ohmori T, Tsuchiya N, Noguchi M, Yuzurihara M, Kase Y, Irahara M: **Changes in circulating cytokine levels in midlife women with psychological symptoms with selective serotonin reuptake inhibitor and Japanese traditional medicine.** *Maturitas* 2009, **62**:146–152.
48. Toriizuka K, Kamiki H, Ohmura NY, Fujii M, Hori Y, Fukumura M, Hirai Y, Isoda S, Nemoto Y, Ida Y: **Anxiolytic effect of Gardeniae Fructus-extract containing active ingredient from Kamishoyosan (KSS), a Japanese traditional Kampo medicine.** *Life Sci* 2005, **77**:3010–3020.
49. Park HJ, Shim HS, Chung SY, Lee TH, Shim I: **Soyo-san reduces depressive-like behavior and proinflammatory cytokines in ovariectomized female rats.** *BMC Complement Altern Med* 2014, **14**:34.
50. Xu F, Peng D, Tao C, Yin D, Kou J, Zhu D, Yu B: **Anti-depression effects of Danggui-Shaoyao-San, a fixed combination of Traditional Chinese Medicine, on depression model in mice and rats.** *Phytomedicine* 2011, **18**:1130–1136.
51. Stefek M, Benes L: **In vitro studies on the activity of Japanese kampo herbal medicines Oren-Gedoku-To (TJ-15) and Toki-Shakuyaku-San (TJ-23) as scavengers of free radicals.** *Drug Metabol Drug Interact* 1994, **11**:25–36.
52. Koike K, Ohno S, Takahashi N, Suzuki N, Nozaki N, Murakami K, Sugiura K, Yamada K, Inoue M: **Efficacy of the herbal medicine Unkei-to as an adjunctive treatment to hormone replacement therapy for postmenopausal women with depressive symptoms.** *Clin Neuropharmacol* 2004, **27**:157–162.
53. Duvan CI, Cumaoglu A, Turhan NO, Karasu C, Kafali H: **Oxidant/antioxidant status in premenstrual syndrome.** *Arch Gynecol Obstet* 2011, **283**:299–304.
54. Scheid V: **Traditional Chinese medicine—what are we investigating? The case of menopause.** *Complement Ther Med* 2007, **15**:54–68.
55. Chini F, Pezzotti P, Orzella L, Borgia P, Guasticchi G: **Can we use the pharmacy data to estimate the prevalence of chronic conditions? a comparison of multiple data sources.** *BMC Public Health* 2011, **11**:688.
56. Zhang LD, Zhang YL, Xu SH, Zhou G, Jin SB: **Traditional Chinese medicine typing of affective disorders and treatment.** *Am J Chin Med* 1994, **22**:321–327.
57. Scheid V: **Depression, constraint, and the liver: (dis)assembling the treatment of emotion-related disorders in Chinese medicine.** *Cult Med Psychiatry* 2013, **37**:30–58.
58. Scheid V, Ward T, Tuffrey V: **Comparing TCM textbook descriptions of menopausal syndrome with the lived experience of London women at midlife and the implications for Chinese medicine research.** *Maturitas* 2010, **66**:408–416.

doi:10.1186/1472-6882-14-206

Cite this article as: Chen et al.: Identifying Chinese herbal medicine for premenstrual syndrome: implications from a nationwide database. *BMC Complementary and Alternative Medicine* 2014 **14**:206.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

